

# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
12850 Memorial Drive  
Houston, TX 77024  
(713)722-7247

**PATIENT** NELVA BRUNSTING      **DOCTOR** A. JAIN, MD      **DRUG DESCRIPTION**  
**BIRTH DATE** 10/08/26  
**MEDICATION** SPIRIVA CAPS 30'S & HANDIHALER  
**QUANTITY** 30      **PATIENT ALLERGIES**  
**DIRECTIONS** INHALE CONTENTS OF ONE CAPSULE ONCE DAILY USING HANDIHALER

*G Losen*

**INGREDIENT NAME:** TIOTROPIUM BROMIDE (TYE-oh-TROE-pee-um BROE-mide)

**COMMON USES:** This medicine is an anticholinergic agent used to treat the symptoms of chronic obstructive pulmonary disease, including chronic bronchitis and emphysema. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking another anticholinergic (eg, ipratropium). Ask your doctor if you are unsure if any of your medicines are anticholinergics. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including if you have trouble urinating; an enlarged prostate; bladder blockage; glaucoma; kidney problems; allergy to milk proteins; pregnancy; or breast-feeding. Tell your doctor if you are having an asthma attack or increased difficulty breathing. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. An additional patient information leaflet is available with this medicine. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have before using this medicine. **DO NOT SWALLOW THE CAPSULES. THE CAPSULES ARE USED WITH A SPECIAL INHALER.** **DO NOT** remove a capsule from the packaging until you are ready to use it. To remove a capsule from the packaging, carefully peel the foil back to expose the capsule, then tip the capsule out of the blister. **DO NOT** cut the foil or use sharp objects to remove the capsule from the blister. If a second capsule is exposed to the air when you are removing a capsule for use, it must be discarded. Do not save the capsule for later. Place the capsule in the special inhaler device immediately. The device will puncture the capsule so that the medicine inside may be inhaled into the lungs through the mouthpiece. Exhale slowly and deeply. **DO NOT** breathe into the mouthpiece of the inhaler. Position the inhaler mouthpiece between your lips and try to rest your tongue flat. Keep your head upright. Do not block the air vents on the inhaler. **TAKE A SLOW, DEEP BREATH.** You should hear or feel the capsule vibrate inside the inhaler. Hold your breath as long as it feels comfortable, then exhale slowly through pursed lips. Breathe out completely. To be sure all of the medicine has been inhaled from the capsule, inhale from the mouthpiece a second time following the same process. **ASK YOUR DOCTOR OR PHARMACIST** if you are unclear on how to use this device or inhale the medicine. **CLEAN THE INHALER DEVICE** once a month according to the instructions in the patient leaflet. Each inhaler device may be used for up to 1 year and then should be replaced. **STORE THIS MEDICINE** at room temperature at 77 degrees F (25 degrees C), away from heat, moisture, and light. Brief storage between 59 and 86 degrees F (15 and 30 degrees C) is permitted. Do not store in the bathroom. This medicine will work best if used regularly. Taking your dose at the same time each day will help you to remember. **IF YOU MISS A DOSE OF THIS MEDICINE,** use it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **DO NOT** use 2 doses at once. **DO NOT** use this medicine more often than 1 time every 24 hours.

**CAUTIONS:** **DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine or to atropine or related medicines (eg, ipratropium). **THIS MEDICINE WILL NOT STOP AN ASTHMA ATTACK** once one has started. **IF YOU ARE ALSO USING A RESCUE INHALER** (eg, albuterol), be sure to always carry the bronchodilator inhaler with you to use during asthma attacks. **DO NOT EXCEED THE RECOMMENDED DOSE** without checking with your doctor. **DO NOT STOP USING THIS MEDICINE** without first checking with your doctor. This medicine may cause dizziness or blurred vision. These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. **AVOID GETTING THIS MEDICINE IN YOUR EYES.** If you get the medicine in your eyes and eye pain, blurred vision, or other vision changes occur, contact your doctor immediately. **THIS MEDICINE MAY SOMETIMES CAUSE SEVERE BREATHING PROBLEMS** right after you use a dose. If this happens, use your short-acting bronchodilator inhaler. Contact your doctor or seek other medical care at once. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE,** either prescription or over-the-counter, check with your doctor or pharmacist. **USE THIS MEDICINE WITH CAUTION IN THE ELDERLY;** they may be more sensitive to its effects, especially constipation and urinary tract infections. **FOR WOMEN:** If you

**PLAN ON BECOMING PREGNANT,** discuss with your doctor the benefits and risks of using this medicine during pregnancy. **IT IS NOT KNOWN IF THIS MEDICINE IS FOUND** in breast milk. **IF YOU ARE OR WILL BE BREAST-FEEDING** while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** **SIDE EFFECTS** that may occur while using this medicine include dry mouth, constipation, sinus inflammation, throat irritation, stomach pain, vomiting, blurred vision, or mild nosebleed. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience irritation, pain, or white patches in your mouth or on your tongue; mouth sores; severe or persistent nosebleeds; severe or persistent constipation; difficult or painful urination; eye pain or discomfort; vision changes (eg, halos, colored images); chest pain; fast or irregular heartbeat; difficulty swallowing; new or worsened breathing problems; or wheezing. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** **IF OVERDOSE IS SUSPECTED,** contact your local poison control center or emergency room immediately. Symptoms may include mental changes; severe constipation; stomach pain; or tremors.

**ADDITIONAL INFORMATION:** **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** obtain refills before your supply runs out.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

**KEEP OUT OF REACH OF CHILDREN. STORE IN SAFETY CONTAINER OR SECURE AREA.**

**NELVA BRUNSTING**

13630 Pinrock, Houston, TX 770797517  
(713)464-4391

**RX # 1496586-03328**      **DATE:** 12/05/10

**SPIRIVA CAPS 30'S & HANDIHALER**

**QTY:** 30      **3 REFILLS BEFORE 12/05/11**  
**New**      **NDC:** 00597-0075-41  
**Retail Price:** \$236.99      **Your Insurance Saved You:** \$215.99

**\$ 21.00**

A. JAIN, MD  
MFG:BOEHRINGER  
TPL/RJW/RJW/RJW/RJW

PLAN: PAIDMPD  
GROUP# CMD3896  
CLAIM REF# RDC9TAQ

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Customer Receipt

**NELVA BRUNSTING**

13630 Pinrock, Houston, TX 770797517  
(713)464-4391

**RX # 1496586-03328**      **DATE:** 12/05/10

**SPIRIVA CAPS 30'S & HANDIHALER**

**QTY:** 30      **3 REFILLS BEFORE 12/05/11**  
**New**      **NDC:** 00597-0075-41  
**Retail Price:** \$236.99      **Your Insurance Saved You:** \$215.99

**\$ 21.00**

A. JAIN, MD  
MFG:BOEHRINGER  
TPL/RJW/RJW/RJW/RJW

PLAN: PAIDMPD  
GROUP# CMD3896  
CLAIM REF# RDC9TAQ

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

**SUN** 1:29PM      **SPIRIVA CAPS 30'S & HANDIHALER**      **QTY 30**  
**New**      **ALPHA**      **00597-0075-41**

TPL/RJW/RJW/RJW/RJW

Brunsting004489

AABrunsting.Financials003940

Get a **FREE** one-on-one  
**Medicare Part D**  
review session!

Your pharmacist will:

- Review your medications and look for ways to help save you money
- Talk about other things you can do to stay well
- Check for potential drug interactions



*Make your appointment today!*

**Walgreens**

There's a way to stay well.

Information provided by Walgreens is not a substitute for your own research concerning your healthcare, including Medicare.



Don't take chances with your health:

**Fill ALL your prescriptions at one pharmacy.**

At Walgreens we'll automatically screen for drug interactions and may be able to offer advice on money-saving alternatives.

For details, speak to our pharmacy staff.

**Walgreens**

There's a way to stay well.



NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

• Your Insurance Saved You: \$215.99

SUN 1:29PM  
\$21.00  
EXPRESS PAY

12/05/10  
New



**Personal Prescription Information**

**LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.**

Take advantage of these convenient services:

- **Auto Refills**  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.
- **Touch Tone Refills**  
Save time by using our automated system for a refill. Just dial the number on your prescription label.

**Walgreens**

The Pharmacy America Trusts • Since 1901™


Visit us online at [Walgreens.com](http://Walgreens.com)

***Thank you for choosing Walgreens!***

Brunsting004490

# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
12850 Memorial Drive  
Houston, TX 77024  
(713)722-7247

<b>PATIENT</b>	NELVA BRUNSTING	<b>DOCTOR</b>	A. JAIN, MD	<b>DRUG DESCRIPTION</b>	 PINK FRONT: m 32
<b>BIRTH DATE</b>	10/08/26	<b>PATIENT ALLERGIES</b>			
<b>MEDICATION</b>	METOPROLOL TARTRATE 50MG TABLETS				
<b>QUANTITY</b>	60				
<b>DIRECTIONS</b>	TAKE 1 TABLET BY MOUTH EVERY 12 HOURS				
	9 am 3 9 pm				

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** METOPROLOL (me-TOE-proe-lole)

**COMMON USES:** This medicine is a beta-adrenergic blocking agent (beta-blocker) used to treat high blood pressure. It may be used alone or with other medicines. It is also used for the long-term treatment of chest pain (angina) and to reduce the risk of death due to heart problems in certain patients who have had a heart attack. It may also be used for other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** **WARNING:** Do not suddenly stop taking this medicine; sharp chest pain, irregular heartbeat, and sometimes heart attack may occur. The risk may be greater if you have certain types of heart disease. Your doctor should slowly lower your dose over several weeks if you need to stop taking it, even if you only take it for high blood pressure. Heart disease is common and you may not know you have it. If you have chest pain or other heart problems occur, contact your doctor right away. You may need to start taking this medicine again. Tell your doctor or dentist that you take this medicine before you receive any medical or dental care, emergency care, or surgery. Tell your doctor if you have a history of diabetes or take medicine to lower your blood sugar (eg, glyburide, insulin). This medicine may hide signs of low blood sugar such as fast heartbeat. Tell your doctor right away if you notice other signs of low blood sugar such as anxiety, chills, dizziness, drowsiness, fainting, headache, tremor, unusual sweating, vision changes, or weakness. This medicine should not usually be used by patients who have a history of certain lung or breathing problems (eg, asthma) or a certain type of adrenal gland tumor (pheochromocytoma). It may worsen these conditions. Tell your doctor if you have a history of these conditions; your doctor may need to adjust your dose or prescribe additional medicine to reduce the risk of side effects. Tell your doctor if you have a history of overactive thyroid. This medicine may hide symptoms of overactive thyroid (eg, fast heartbeat). Do not suddenly stop taking this medicine; suddenly stopping this medicine could worsen your condition. Your doctor should slowly lower your dose over several weeks if you need to stop taking it. Check with your doctor for more information. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking mifepristone. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking amiodarone, bupropion, certain HIV protease inhibitors (eg, ritonavir), certain selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), cimetidine, digoxin, diphendramine, disopyramide, flecainide, hormonal contraceptives (eg, birth control pills), hydralazine, hydroxychloroquine, ketanserin, mefloquine, phenothiazines (eg, thioridazine), propafenone, quinazoline (eg, aluzosin), quinidine, terbinafine, thiazines (eg, propylthiouracil, verapamil), barbiturates (eg, phenobarbital), indomethacin, phenylpropanolamine, bupivacaine, lidocaine, or clonidine. **DO NOT STOP** or change your medicine without your doctor or pharmacist approval. Inform your doctor of any other medical conditions including lung or breathing problems (eg, asthma, bronchitis, chronic obstructive pulmonary disease (COPD)), diabetes, low blood pressure, thyroid disease, or adrenal gland tumor (pheochromocytoma), allergies, pregnancy, or breastfeeding. Tell your doctor if you have a history of heart attack, slow or irregular heartbeat, heart failure, or other heart problems; chest pain or angina; blood vessel or circulation problems; liver disease; or if you will be having surgery or receiving anesthesia. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have a very slow heartbeat (eg, bradycardia), certain types of irregular heartbeat (eg, atrioventricular [AV] block, sick sinus syndrome), moderate to severe heart failure, very low systolic blood pressure (less than 100 mm Hg), or severe blood circulation problems. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. **TAKE THIS MEDICINE BY MOUTH WITH FOOD** or immediately following meal at the same time each day. **STORE THIS MEDICINE** at room temperature, between 59 and 86 degrees F (15 and 30 degrees C). Store away from heat, moisture, and light. Do not store in the bathroom. **IF YOU MISS A DOSE OF THIS MEDICINE,** take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** **DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine or to another beta-blocker (eg, propranolol). **IF YOU HAVE A HISTORY OF ANY SEVERE ALLERGIC REACTION,** talk with your doctor. You may be at risk for an even more severe allergic reaction if you come into contact with the substance that caused your allergy. Some medicines used to treat

severe allergies may also not work as well while you are using this medicine. **LAB TESTS,** including liver and kidney function, blood pressure, and complete blood cell counts, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to keep all doctor and lab appointments. **THIS MEDICINE MAY CAUSE DROWSINESS, DIZZINESS, OR LIGHTHEADEDNESS.** These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. Do not drive or perform other possibly unsafe tasks until you know how you react to it. This medicine may cause dizziness, lightheadedness, or fainting; alcohol, hot weather, exercise, or fever may increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of any of these effects. **PATIENTS WHO TAKE MEDICINE FOR HIGH BLOOD PRESSURE** often feel tired or run down for a few weeks after starting treatment. Be sure to take your medicine even if you may not feel "normal." Tell your doctor if you develop any new symptoms. **DO NOT SUDDENLY STOP TAKING THIS MEDICINE** without first consulting your doctor. If your doctor decides you should no longer use this medicine, you will need to stop this medicine gradually according to your doctor's instructions. **IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK YOUR BLOOD PRESSURE** and heart rate regularly, be sure to do so. **TELL YOUR DOCTOR OR DENTIST** that you take this medicine before you receive any medical or dental care, emergency care, or surgery. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescription or over-the-counter, check with your doctor or pharmacist. Do not take any medicines used to treat colds or congestion without first talking with your doctor or pharmacist. **FOR WOMEN:** If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. This medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby. **DIABETES PATIENTS:** this medicine may hide signs of low blood sugar, such as rapid heartbeat. Be sure to watch for other signs or low blood sugar. Low blood sugar may make you anxious, sweaty, weak, dizzy, drowsy, or faint. It may also make your vision change; give you a headache, chills, or tremors; or make you more hungry. Check blood sugar levels closely. Ask your doctor before you change the dose of your diabetes medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include constipation, diarrhea, dizziness, dry mouth/eyes, gas, headache, heartburn, lightheadedness, mild drowsiness, muscle aches, nausea, stomach pain, trouble sleeping, unusual tiredness or weakness, or vomiting. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience blue or unusually cold hands or feet; chest pain; fainting; hallucinations; mood or mental changes (eg, confusion, depression); pounding in the chest; severe dizziness; or lightheadedness; shortness of breath; slow or irregular heartbeat; swelling of the arms, hands, and feet; vision changes; wheezing; yellowing of the skin or eyes. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include chest pain, seizures, very slow heart beat, severe dizziness, fainting, and difficult or slowed breathing.

**ADDITIONAL INFORMATION:** **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** obtain refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.


Do not flush unused medications or pour down a sink or drain.

WIC# 957918

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

<p><b>NELVA BRUNSTING</b> 13630 Pinerock, Houston, TX 770797517 (713)948-4391 <b>RX # 1496588-03328</b>      DATE: 12/05/10</p>	<p><b>NELVA BRUNSTING</b> 13630 Pinerock, Houston, TX 770797517 (713)948-4391 <b>RX # 1496588-03328</b>      DATE: 12/05/10</p>
<p><b>METOPROLOL TARTRATE 50MG TABLETS</b> QTY: 60    3 REFILLS BEFORE 12/05/11 New    NDC:00378-0032-10 Retail Price: \$14.99    Your Insurance Saved You: \$10.61    <b>\$ 4.38</b></p>	<p><b>METOPROLOL TARTRATE 50MG TABLETS</b> QTY: 60    3 REFILLS BEFORE 12/05/11 New    NDC:00378-0032-10 Retail Price: \$14.99    Your Insurance Saved You: \$10.61    <b>\$ 4.38</b></p>
<p>A. JAIN, MD MFG:MYLAN TPL/RJW/RJW /RJW</p>	<p>A. JAIN, MD MFG:MYLAN TPL/RJW/RJW /RJW</p>
<p>PLAN: PAIDMPD GROUP# CMD3896 CLAIM REF# YRKMEWD</p>	<p>PLAN: PAIDMPD GROUP# CMD3896 CLAIM REF# YRKMEWD</p>
<p><b>Walgreens</b>    12850 MEMORIAL DRIVE HOUSTON, TX 77024 PH: (713)722-7247    Customer Receipt</p>	<p><b>Walgreens</b>    12850 MEMORIAL DRIVE HOUSTON, TX 77024 PH: (713)722-7247    Duplicate Receipt</p>

Pharmacy use only

<p>SUN 1:29PM New</p>	<p>METOPROLOL TARTRATE 50MG TABLETS 00378-0032-10 CELL 138</p>	<p>QTY 60 20 DRAM</p>	<p> PINK FRONT: m 32</p>
---------------------------	--	---------------------------	---

TPLRJW/RJW /RJW

Brunsting004481

## Educate Yourself About Stroke

Stroke kills more women than breast cancer, cervical cancer, uterine cancer, and ovarian cancer combined<sup>1</sup>

### Be Prepared...

#### ACT F.A.S.T!

#### Stroke risk factors

<b>FACE</b>	Facial droop Uneven smile	High blood pressure High cholesterol Heart disease
<b>ARM</b>	Arm numbness Arm weakness	Diabetes Smoking
<b>SPEECH</b>	Slurred speech Difficulty speaking or understanding	Heavy alcohol use Physical inactivity and obesity
<b>TIME</b>	Call 911 and get to the hospital immediately.	Atrial fibrillation (irregular heartbeat) Family history of stroke

#### Check out National Stroke Association's Virtual Health Fair, January 19, 2011 and beyond

- Register now and get comprehensive, accurate, and up-to-date education on and resources for stroke prevention and recovery
- Visit [www.stroke.org/catalina](http://www.stroke.org/catalina), hosted by National Stroke Association

For more information on stroke,  
visit [www.getstrokeinfo.com](http://www.getstrokeinfo.com)

Reference: 1. Heron M, Hoyert DL, Murphy SL, Xu J, Kochanek KD, Tejada-Vera B. Deaths: final data for 2008. *Natl Vital Stat Rep*. 2009;57(14):1-134.

Brought to you by  
**Genentech**  
A Member of the Roche Group

©2010 Genentech USA, Inc. All rights reserved. AC1000107200 Printed in USA.

You're taking action  
*by managing the most important part of your life, your health.*



Having your prescriptions refilled on time is a positive step towards managing your condition, but it's just one part of managing it. Eating healthy foods and getting proper exercise are also important factors for total health. If you don't already have a diet and exercise regimen, then talk to your doctor about the best plan for you.

You are taking **metoprolol** to help control your high blood pressure. Below are some key points to remember about **metoprolol**.

- Follow the directions on your prescription bottle carefully.
- Take your medicine exactly as your doctor prescribed.
- Continue to take it even if you feel well. Give your body time to adjust to the medicine.
- To help you remember, take it around the same time(s) every day.
- Check your blood pressure regularly.
- If you have any questions, ask one of our pharmacists on the pharmacy team or call your doctor. It's important for you to keep all scheduled appointments for lab testing.
- Talk to your doctor about how you feel and about any symptoms you have during the starting of this medicine.

The information above is brought to you as a courtesy from your pharmacist.

8  
BR




SUN 1:29PM  
\$4.38  
EXPRESS PAY

NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

12/05/10  
New

• Your Insurance Saved You: \$10.61



## Personal Prescription Information

#### LOOK INSIDE FOR IMPORTANT INFORMATION

##### ABOUT YOUR MEDICATION.

##### Take advantage of these convenient services:

- **Touch Tone Refills**  
Save time by using our automated system for a refill. Just dial the number on your prescription label.
- **Auto Refills**  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

**Walgreens**  
The Pharmacy America Trusts • Since 1901™  
Visit us online at [Walgreens.com](http://Walgreens.com)

**Thank you for choosing Walgreens!**

To stop receiving sponsored info at this pharmacy, call 888-336-5744 or visit [remove-me.net](http://remove-me.net) Use code:2875677 339 0223746

Brunsting004482

AA Brunsting.Financials003942


OIRT: 2875 677 339 0223746

INFO: 0896 00378003210

3440

# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
12850 Memorial Drive  
Houston, TX 77024  
(713)722-7247

**PATIENT** NELVA BRUNSTING **DOCTOR** A. JAIN, MD **DRUG DESCRIPTION**  
**BIRTH DATE** 10/08/26  
**MEDICATION** METOPROLOL TARTRATE 50MG TABLETS **PATIENT ALLERGIES**  
**QUANTITY** 60  
**DIRECTIONS** TAKE 1 TABLET BY MOUTH EVERY 12 HOURS  
  
**PINK FRONT: m 32**

**INGREDIENT NAME:** METOPROLOL (me-TOE-proe-loe)

**COMMON USES:** This medicine is a beta-adrenergic blocking agent (beta-blocker) used to treat high blood pressure. It may be used alone or with other medicines. It is also used for the long-term treatment of chest pain (angina) and to reduce the risk of death due to heart problems in certain patients who have had a heart attack. It may also be used for other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE: WARNING:** Do not suddenly stop taking this medicine; sharp chest pain, irregular heartbeat, and sometimes heart attack may occur. The risk may be greater if you have certain types of heart disease. Your doctor should slowly lower your dose over several weeks if you need to stop taking it, even if you only take it for high blood pressure. Heart disease is common and you may not know you have it. Limit physical activity while you are lowering your dose. If new or worsened chest pain or other heart problems occur, contact your doctor right away. You may need to start taking this medicine again. Tell your doctor or dentist that you take this medicine before you receive any medical or dental care, emergency care, or surgery. Tell your doctor if you have a history of diabetes or take medicine to lower your blood sugar (eg, glyburide, insulin). This medicine may hide signs of low blood sugar such as fast heartbeat. Tell your doctor right away if you notice other signs of low blood sugar such as anxiety, chills, dizziness, drowsiness, fainting, headache, tremor, unusual sweating, vision changes, or weakness. This medicine should not usually be used by patients who have a history of certain lung or breathing problems (eg, asthma) or a certain type of adrenal gland tumor (pheochromocytoma). It may worsen these conditions. Tell your doctor if you have a history of these conditions; your doctor may need to adjust your dose or prescribe additional medicine to reduce the risk of side effects. Tell your doctor if you have a history of overactive thyroid. This medicine may hide symptoms of overactive thyroid (eg, fast heartbeat). Do not suddenly stop taking this medicine; suddenly stopping this medicine could worsen your condition. Your doctor should slowly lower your dose over several weeks if you need to stop taking it. Check with your doctor for more information. Other medicines or medical conditions may interact with this medicine.

**INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking mifepridil. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking amiodarone, bupropion, certain HIV protease inhibitors (eg, ritonavir), certain selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine, cimetidine, digoxin, trimethoprim, disopyramide, flecainide, hormonal contraceptives (eg, birth control pills), hydroxyzine, hydroxychloroquine, ketanserin, mefloquine, phenothiazines (eg, thioridazine), propafenone, quinazolinones (eg, alfluzonil), quinidine, terbinafine, thiamines (eg, propylthiouracil), verapamil, barbiturates (eg, phenobarbital), indometacin, phenylpropranolamine, bupivacaine, lidocaine, or clonidine. **DO NOT STOP OR STOP ANY MEDICINE** without doctor or pharmacist approval. Inform your doctor of any other medical conditions including lung or breathing problems (eg, asthma, bronchitis, chronic obstructive pulmonary disease [COPD]), diabetes, low blood pressure, thyroid problems, or adrenal gland tumor (pheochromocytoma), allergies, pregnancy, or breastfeeding. Tell your doctor if you have a history of heart attack, slow or irregular heartbeat, heart failure, or other heart problems; chest pain or angina; blood vessel or circulation problems; liver disease; or if you will be having surgery or receiving anesthesia. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have a very slow heartbeat (eg, bradycardia), certain types of irregular heartbeat (eg, atrioventricular (AV) block, sick sinus syndrome), moderate to severe heart failure, very low systolic blood pressure (less than 100 mm Hg), or severe blood circulation problems. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. TAKE THIS MEDICINE BY MOUTH WITH FOOD or immediately following a meal at the same time each day. STORE THIS MEDICINE at room temperature, between 59 and 86 degrees F (15 and 30 degrees C). Store away from heat, moisture, and light. Do not store in the bathroom. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** DO NOT USE THIS MEDICINE if you are allergic to any ingredient in this medicine or to another beta-blocker (eg, propranolol). IF YOU HAVE A HISTORY OF ANY SEVERE ALLERGIC REACTION, talk with your doctor. You may be at risk for an even more severe allergic reaction if you come into contact with the substance that caused your allergy. Some medicines used to treat

severe allergies may also not work as well while you are using this medicine. LAB TESTS, including liver and kidney function, blood pressure, and complete blood cell counts, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to keep all doctor and lab appointments. THIS MEDICINE MAY CAUSE DROWSINESS, DIZZINESS, OR LIGHTHEADEDNESS. These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. Do not drive or perform other possibly unsafe tasks until you know how you react to it. This medicine may cause dizziness, lightheadedness, or fainting; alcohol, hot weather, exercise, or fever may increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of any of these effects. PATIENTS WHO TAKE MEDICINE FOR HIGH BLOOD PRESSURE often feel tired or run down for a few weeks after starting treatment. Be sure to take your medicine even if you may not feel "normal." Tell your doctor if you develop any new symptoms. **DO NOT SUDDENLY STOP TAKING THIS MEDICINE** without first consulting your doctor. If your doctor decides you should no longer use this medicine, you will need to stop this medicine gradually according to your doctor's instructions. IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK YOUR BLOOD PRESSURE and heart rate regularly, be sure to do so. TELL YOUR DOCTOR OR DENTIST that you take this medicine before you receive any medical or dental care, emergency care, or surgery. BEFORE YOU BEGIN TAKING ANY NEW MEDICINES, either prescription or over-the-counter, check with your doctor or pharmacist. Do not take any medicines used to treat colds or congestion without first talking with your doctor or pharmacist. FOR WOMEN: If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. This medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby. **DIABETES PATIENTS:** this medicine may hide signs of low blood sugar, such as rapid heartbeat. Be sure to watch for other signs of low blood sugar. Low blood sugar may make you anxious, sweaty, weak, dizzy, drowsy, or faint. It may also make your vision change; give you a headache, chills, or tremors; or make you more hungry. Check blood sugar levels closely. Ask your doctor before you change the dose of your diabetes medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include constipation, diarrhea, dizziness, dry mouth/eyes, gas, headache, heartburn, lightheadedness, mild drowsiness, muscle aches, nausea, stomach pain, trouble sleeping, yellowing of the skin or eyes, weight loss, or vomiting. If any of these bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience blue or unusually cold hands or feet; chest pain; fainting; hallucinations; mood or mental changes (eg, confusion, depression); pounding in the chest; severe dizziness or lightheadedness; shortness of breath; slow or irregular heartbeat; swelling of the arms, hands, and feet; vision changes; wheezing; yellowing of the skin or eyes. AN ALLERGIC REACTION to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include chest pain, seizures, very slow heart beat, severe dizziness, fainting, and difficult or slowed breathing.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

NELVA BRUNSTING

13830 Pinerock, Houston, TX 770797517  
(713)464-4391

RX # 1496588-03328

DATE: 01/01/11

METOPROLOL TARTRATE 50MG TABLETS

QTY: 60 2 REFILLS BEFORE 12/05/11

Refill NDC: 00378-0032-10

Retail Price: \$14.99 Your Insurance Saved You: \$10.61

\$ 4.38

A. JAIN, MD PLAN: PAID/MD  
MFG: MYLAN GROUP# CMD3896  
XXX/JC/JC/ /RJW CLAIM REF# YXEP3F1

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Customer Receipt

Pharmacy-use only

SUN 12:00PM  
Refill

METOPROLOL TARTRATE 50MG TABLETS  
00378-0032-10  
CELL 138

NELVA BRUNSTING

13830 Pinerock, Houston, TX 770797517  
(713)464-4391

RX # 1496588-03328

DATE: 01/01/11

METOPROLOL TARTRATE 50MG TABLETS

QTY: 60 2 REFILLS BEFORE 12/05/11

Refill NDC: 00378-0032-10

Retail Price: \$14.99 Your Insurance Saved You: \$10.61

\$ 4.38

A. JAIN, MD PLAN: PAID/MD  
MFG: MYLAN GROUP# CMD3896  
XXX/JC/JC/ /RJW CLAIM REF# YXEP3F1

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Duplicate Receipt



PINK  
FRONT: m 32

XXX/JC/JC/ /RJW

Brunsting004479

AABrunsting.Financials003944

## Educate Yourself About Stroke

Stroke kills more women than breast cancer, cervical cancer, uterine cancer, and ovarian cancer combined<sup>1</sup>

### Be Prepared...

ACT F.A.S.T!		Stroke risk factors
<b>F</b> ACE	Facial droop Uneven smile	High blood pressure High cholesterol Heart disease
<b>A</b> RM	Arm numbness Arm weakness	Diabetes Smoking Heavy alcohol use
<b>S</b> PEECH	Slurred speech Difficulty speaking or understanding	Physical inactivity and obesity Atrial fibrillation (irregular heartbeat)
<b>T</b> IME	Call 911 and get to the hospital immediately.	Family history of stroke

### Check out National Stroke Association's Virtual Health Fair, January 19, 2011 and beyond

- Register now and get comprehensive, accurate, and up-to-date education on and resources for stroke prevention and recovery
- Visit [www.stroke.org/catalina](http://www.stroke.org/catalina), hosted by National Stroke Association

For more information on stroke,  
visit [www.getstrokeinfo.com](http://www.getstrokeinfo.com)

Reference: 1. Heron M, Hoyert DL, Murphy SL, Xu J, Kochanek KD, Tejada-Vera B. Deaths: final data for 2008. *Natl Vital Stat Rep*. 2008;57(14):1-134.

Brought to you by **Genentech**  
A Member of the Roche Group

©2010 Genentech USA, Inc. All rights reserved. AC10800107200 Printed in USA.

You're taking action  
by managing the most important part of your life, your health.



Having your prescriptions refilled on time is a positive step towards managing your condition, but it's just one part of managing it. Eating healthy foods and getting proper exercise are also important factors for total health. If you don't already have a diet and exercise regimen, then talk to your doctor about the best plan for you.

You are taking **metoprolol** to help control your high blood pressure. Below are some key points to remember about **metoprolol**.

- Follow the directions on your prescription bottle carefully.
- Take your medicine exactly as your doctor prescribed.
- Continue to take it even if you feel well. Give your body time to adjust to the medicine.
- To help you remember, take it around the same time(s) every day.
- Check your blood pressure regularly.
- If you have any questions, ask one of our pharmacists on the pharmacy team or call your doctor. It's important for you to keep all scheduled appointments for lab testing.
- Talk to your doctor about how you feel and about any symptoms you have during the starting of this medicine.

The information above is brought to you as a courtesy from your pharmacist.




SUN 12:00PM  
\$4.38  
EXPRESS PAY

NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

01/01/11  
Refill

- Store TT 1 of 5
- Your Insurance Saved You: \$10.61



## Personal Prescription Information

### LOOK INSIDE FOR IMPORTANT INFORMATION

#### ABOUT YOUR MEDICATION.

#### Take advantage of this convenient service:

- **Auto Refills**  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

**Walgreens**

The Pharmacy America Trusts • Since 1901\*

Visit us online at [Walgreens.com](http://Walgreens.com)


**Thank you for choosing Walgreens!**

To stop receiving sponsored info at this pharmacy, call 888-336-5744 or visit [remove-me.net](http://remove-me.net) Use code:2875677 002 0240051

Brunsting004480

# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
 12850 Memorial Drive  
 Houston, TX 77024  
 (713)722-7247

<b>PATIENT</b> NELVA BRUNSTING	<b>DOCTOR</b> A. JAIN, MD	<b>DRUG DESCRIPTION</b>
<b>BIRTH DATE</b> 10/08/26		 WHITE FRONT: L U BACK: C32
<b>MEDICATION</b> ETHAMBUTOL 400MG TABLETS	<b>PATIENT ALLERGIES</b>	
<b>QUANTITY</b> 90		
<b>DIRECTIONS</b> TAKE 3 TABLETS BY MOUTH EVERY DAY		

**INGREDIENT NAME:** ETHAMBUTOL (e-THAM-byoo-tole)

**COMMON USES:** This medicine is an antibacterial used to treat tuberculosis (TB).

**BEFORE USING THIS MEDICINE:** INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. Inform your doctor of any other medical conditions, allergies, pregnancy, or breast-feeding.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. **THIS MEDICINE MAY BE TAKEN WITH FOOD** if it upsets your stomach. **STORE THIS MEDICINE** at room temperature in a tightly-closed container, away from heat and light. **IF YOU MISS A DOSE OF THIS MEDICINE**, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** DO NOT STOP USING THIS MEDICINE without first checking with your doctor. **THIS MEDICINE MAY CAUSE dizziness.** Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. **HAVE REGULAR EYE EXAMINATIONS** while you are taking this medicine even if you do not notice changes in your vision. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE**, either prescription or over-the-counter, check with your doctor or pharmacist.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS, that may go away during

treatment, include nausea, headache, or dizziness. If they continue or are bothersome, check with your doctor. **CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE** if you experience changes in vision, vomiting, skin rash, itching, fever, confusion, or numbness or tingling of extremities. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately.

**ADDITIONAL INFORMATION:** If your symptoms do not improve within 2 to 3 weeks, or if they become worse, check with your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME**, obtain refills before your supply runs out.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

**NELVA BRUNSTING**  
 13630 Pinerock, Houston, TX 770797517  
 (713)464-4391  
**RX # 1494792-03328**      DATE: 01/01/11

**ETHAMBUTOL 400MG TABLETS**  
 QTY: 90      2 REFILLS BEFORE 11/29/11  
 Refill      NDC: 68180-0281-01  
 Retail Price: \$153.59      Your Insurance Saved You: \$148.59      \$ 5.00

A. JAIN, MD      PLAN: PAIDMPP  
 MFG: LUPIN      GROUP# CMD3596  
 XXX/JIC/JIC /RJW      CLAIM REF# SGA93NQ

**Walgreens**      12850 MEMORIAL DRIVE HOUSTON, TX 77024  
 PH: (713)722-7247

Customer Receipt

**NELVA BRUNSTING**  
 13630 Pinerock, Houston, TX 770797517  
 (713)464-4391  
**RX # 1494792-03328**      DATE: 01/01/11

**ETHAMBUTOL 400MG TABLETS**  
 QTY: 90      2 REFILLS BEFORE 11/29/11  
 Refill      NDC: 68180-0281-01  
 Retail Price: \$153.59      Your Insurance Saved You: \$148.59      \$ 5.00

A. JAIN, MD      PLAN: PAIDMPP  
 MFG: LUPIN      GROUP# CMD3596  
 XXX/JIC/JIC /RJW      CLAIM REF# SGA93NQ

**Walgreens**      12850 MEMORIAL DRIVE HOUSTON, TX 77024  
 PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

SUN 12:00PM  
 Refill

ETHAMBUTOL 400MG TABLETS  
 68180-0281-01  
 ALPHA

QTY 90



WHITE  
 FRONT: L U  
 BACK: C32

XXX/JIC/JIC /RJW

Brunsting004473

Get a **FREE** one-on-one  
**Medicare Part D**  
review session!

Your pharmacist will:

- Review your medications and look for ways to help save you money
- Talk about other things you can do to stay well
- Check for potential drug interactions



*Make your appointment today!*

**Walgreens**

There's a way to stay well.

Information provided by Walgreens is not a substitute for your own research concerning your healthcare, including Medicare.

**Fever at 3 a.m.?**

We have the most 24-hour  
pharmacies in America, so you get  
the help you need, when you need it.



For the 24-hour Walgreens nearest  
you, call 1-800-WALGREENS  
or visit [Walgreens.com/findastore](http://Walgreens.com/findastore).

Another great service from

**Walgreens**

The Pharmacy America Trusts Since 1901

8  
BR



NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

- Store TT 2 of 5
- Your Insurance Saved You: \$148.59

SUN 12:00PM  
**\$5.00**  
EXPRESS PAY

01/01/11  
Refill



**Personal  
Prescription  
Information**

LOOK INSIDE FOR IMPORTANT INFORMATION

**ABOUT YOUR MEDICATION.**

Take advantage of this convenient service:

- **Auto Refills**  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

**Walgreens**

The Pharmacy America Trusts Since 1901

Visit us online at [Walgreens.com](http://Walgreens.com)


**Thank you for choosing Walgreens!**

Brunsting004474



# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
 12850 Memorial Drive  
 Houston, TX 77024  
 (713)722-7247

<b>PATIENT</b> NELVA BRUNSTING	<b>DOCTOR</b> A. JAIN, MD	<b>DRUG DESCRIPTION</b>
<b>BIRTH DATE</b> 10/08/26		 PINK FRONT: 93 BACK: 7146
<b>MEDICATION</b> AZITHROMYCIN 250MG TABLETS	<b>PATIENT ALLERGIES</b>	
<b>QUANTITY</b> 30		
<b>DIRECTIONS</b> TAKE 1 TABLET BY MOUTH EVERY DAY		

**INGREDIENT NAME:** AZITHROMYCIN  
 (ay-ZITH-roe-MYE-sin)

**COMMON USES:** This medicine is a macrolide antibiotic used to treat bacterial infections.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking dofetilide, nitroinib, propafenone, pimozide, or tetrabenazine. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking antiarrhythmics (eg, disopyramide), anticoagulants (eg, warfarin), arsenic, astemizole, carbamazepine, cisapride, digoxin, domperidone, meprobamate, methadone, nelfinavir, cyclosporine, ergot alkaloids (eg, ergotamine), paliperidone, phenytoin, quinolones (eg, levofloxacin), rifampin, terfenadine, theophylline, triazolam, tyrosine kinase inhibitors (eg, dasatinib), or medicines that may affect your heartbeat. Ask your doctor if you are unsure if any of the medicines you are taking may affect your heartbeat. Inform your doctor of any other medical conditions including irregular heartbeat, kidney problems, liver problems, myasthenia gravis, allergies, pregnancy or breastfeeding. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine may be taken on an empty stomach or with food. DO NOT TAKE THIS MEDICINE within 1 hour before or 2 hours after aluminum- or magnesium-containing antacids. STORE THIS MEDICINE at room temperature, away from heat and light. TO CLEAR UP YOUR INFECTION COMPLETELY, continue taking this medicine for the full course of treatment even if you feel better in a few days. Do not miss any doses. Taking this medicine at the same time each day will make it easier to remember. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. If you miss a dose, do not take 2 doses at once.

**CAUTIONS:** DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it, to other macrolide antibiotics (such as erythromycin), to ketolide antibiotics (such as telithromycin), or if you are allergic to any ingredient in this product. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to erythromycin or any macrolide or ketolide antibiotic. A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine, contact your doctor or pharmacist. IF YOU EXPERIENCE difficulty breathing; tightness of chest; swelling of eyelids, face, or lips; or if you develop a rash or hives, tell your doctor immediately. Do not take any more of this medicine unless your doctor tells you to do so. This medicine may cause drowsiness, dizziness, or lightheadedness. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. THIS

MEDICINE MAY CAUSE increased sensitivity to the sun. Avoid exposure to the sun, sunlamps, or tanning booths until you know how you react to this medicine. Use a sunscreen or protective clothing if you must be outside for a prolonged period. MILD DIARRHEA IS COMMON with antibiotic use. However, a more serious form of diarrhea (pseudomembranous colitis) may rarely occur. This may develop while you use the antibiotic or within several months after you stop using it. Contact your doctor right away if stomach pain or cramps, severe diarrhea, or bloody stools occur. Do not treat diarrhea without first checking with your doctor. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. FOR WOMEN: IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. IT IS UNKNOWN IF THIS MEDICINE IS EXCRETED in breast milk. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS, that may go away during treatment, include mild diarrhea, headache, nausea, or stomach pain or upset. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience vomiting, changes in hearing, or ringing in the ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience chest pain; swelling of your hands, legs, face, lips, eyes, throat, or tongue; difficulty swallowing, speaking, or breathing; eye or vision problems; fainting; hearing loss; irregular heartbeat; muscle weakness; reddened, blistered, or swollen skin; seizures; severe stomach pain, cramps, or diarrhea; bloody stools; unusual vaginal itching, odor, or discharge; or yellowing of the eyes or skin. An allergic reaction to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, dizziness, trouble breathing, or unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include nausea, vomiting, stomach upset, and diarrhea.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

**NELVA BRUNSTING**

13630 Pinerock, Houston, TX 770797517  
 (713)464-4391

RX # 1494789-03328

DATE: 01/01/11

**AZITHROMYCIN 250MG TABLETS**

QTY: 30 2 REFILLS BEFORE 11/29/11  
 Refill NDC: 00093-7146-56  
 Retail Price: \$195.79 Your Insurance Saved You: \$190.79

\$ 5.00

A. JAIN, MD  
 MFG: TEVA  
 XXX/KHN/KHN/KHN/RJW

PLAN: PAIDMFD  
 GROUP# CMD3896  
 CLAIM REF# SQA93N1

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
 PH: (713)722-7247

Customer Receipt

**NELVA BRUNSTING**

13630 Pinerock, Houston, TX 770797517  
 (713)464-4391

RX # 1494789-03328

DATE: 01/01/11

**AZITHROMYCIN 250MG TABLETS**

QTY: 30 2 REFILLS BEFORE 11/29/11  
 Refill NDC: 00093-7146-56  
 Retail Price: \$195.79 Your Insurance Saved You: \$190.79

\$ 5.00

A. JAIN, MD  
 MFG: TEVA  
 XXX/KHN/KHN/KHN/RJW

PLAN: PAIDMFD  
 GROUP# CMD3896  
 CLAIM REF# SQA93N1

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
 PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

SUN 12:00PM  
 Refill

AZITHROMYCIN 250MG TABLETS  
 00093-7146-56  
 ALPHA

QTY 30  
 10 DRAM



PINK  
 FRONT: 93  
 BACK: 7146

XXX/KHN/KHN/KHN/RJW

Brunsting004475



## Are you getting the most from your Medicare Part D plan?

We can help you find out with a **FREE one-on-one plan review session** with your pharmacist. We'll review your medications, look for ways to help save you money and more.

*Make your appointment today!*

# Walgreens

There's a way to stay well.

Information provided by Walgreens is not a substitute for your own research concerning your healthcare, including Medicare.

49376

## AARP Members

### Get special offers each week from Walgreens!\*



Just show your AARP card at time of purchase to receive exclusive offers on everyday products - available only to AARP members.



*Smarter. Stronger. Healthier.*

\*AARP card must be presented at time of purchase. For complete program terms and conditions, visit [Walgreens.com/aarpcard](http://Walgreens.com/aarpcard). AARP receives a royalty for the use of its intellectual property. Amounts paid are used for the general purposes of AARP and its members.

DPT: 2875 677 002 0240067 INFO: 0900 00093714556

3438




NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

• Store TT 3 of 5  
• Your Insurance Saved You: \$190.79

SUN 12:00PM  
\$5.00  
EXPRESS PAY

01/01/11  
Refill



## Personal Prescription Information

### LOOK INSIDE FOR IMPORTANT INFORMATION

#### ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- Auto Refills  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

# Walgreens

The Pharmacy America Trusts • Since 1901™


Visit us online at [Walgreens.com](http://Walgreens.com)

Thank you for choosing Walgreens!

Brunsting004476

# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
12850 Memorial Drive  
Houston, TX 77024  
(713)722-7247

<b>PATIENT</b> NELVA BRUNSTING	<b>DOCTOR</b> A. JAIN, MD	<b>DRUG DESCRIPTION</b>
<b>BIRTH DATE</b> 10/08/26		
<b>MEDICATION</b> RIFAMPIN 300MG CAPSULES	<b>PATIENT ALLERGIES</b>	
<b>QUANTITY</b> 60		<b>DARK REDDISH-BROWN</b>
<b>DIRECTIONS</b> TAKE 2 CAPSULES BY MOUTH EVERY DAY		<b>FRONT: LANNETT</b>
		<b>BACK: 1315</b>

**INGREDIENT NAME:** RIFAMPIN (rif-AM-pin)

**COMMON USES:** This medicine is a rifamycin antibiotic used to treat tuberculosis (TB), as well as to treat those who have been exposed to meningitis-causing bacteria before they become sick. This drug does not treat acute meningitis. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking HIV protease inhibitors, pyrazinamide, or birth control pills. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking anticoagulants, cyclosporine, digtoxin, isoniazid, itraconazole, macrolide antibiotics, mexiletine, nevirapine, quinidine, progestins, theophylline, tocinide, verapamil, or medicine for anxiety, sleep, or seizures. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, allergies, pregnancy, or breast-feeding. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. Take this medicine on an empty stomach at least 1 hour before or 2 hours after eating. **THIS MEDICINE MAY BE TAKEN WITH FOOD** if it upsets your stomach. **STORE THIS MEDICINE** at room temperature in a tightly-closed container, away from heat and light. **TO CLEAR UP YOUR INFECTION COMPLETELY**, continue taking this medicine for the full course of treatment. Do not miss any doses. **IF YOU MISS A DOSE OF THIS MEDICINE**, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** IT MAY TAKE SEVERAL WEEKS for this medicine to work. Do not stop using this medicine without checking with your doctor. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. **THIS MEDICINE MAY CAUSE** urine, feces, saliva, sweat, and tears to turn orange or red. **THIS MEDICINE MAY PERMANENTLY STAIN** soft contact lenses. **KEEP ALL DOCTOR AND LABORATORY**

**APPOINTMENTS** while you are using this medicine. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE**, either prescription or over-the-counter, check with your doctor or pharmacist. **USING THIS MEDICINE** while you are taking birth control pills may decrease the effectiveness of your birth control pills. To prevent pregnancy, use an additional form of birth control. **FOR WOMEN: IF YOU PLAN ON BECOMING PREGNANT**, discuss with your doctor the benefits and risks of using this medicine during pregnancy.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS, that may go away during treatment, include stomach upset, heartburn, loss of appetite, nausea, gas, abdominal cramping, headache, drowsiness, dizziness, menstrual changes, joint pain, or leg cramps. If they continue or are bothersome, check with your doctor. **CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE** if you experience vomiting, diarrhea, fever, chills, changes in vision, or confusion. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience severe stomach pain, dark urine, or yellowing of eyes or skin. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include swelling of face or around eyes, itching over the entire body, orange or red discoloration of skin or eyes, nausea, vomiting, drowsiness, and loss of consciousness.

**ADDITIONAL INFORMATION:** **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME**, obtain refills before your supply runs out.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

**NELVA BRUNSTING**  
13630 Finerock, Houston, TX 770797517  
(713)464-4391  
**RX # 1494790-03328**      DATE: 01/01/11

**RIFAMPIN 300MG CAPSULES**  
QTY: 60    2 REFILLS BEFORE 11/29/11  
Refill      NDC:00527-1315-30  
Retail Price: \$113.89    Your Insurance Saved You: \$108.89    \$ 5.00

A: JAIN, MD      PLAN: PAIDMPD  
MFG: LANNETT      GROUP: CMD3896  
XXX/KHN/KHN/KHN/RJW      CLAIM REF# ONA1TWP

**Walgreens**      12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Customer Receipt

**NELVA BRUNSTING**  
13630 Finerock, Houston, TX 770797517  
(713)464-4391  
**RX # 1494790-03328**      DATE: 01/01/11

**RIFAMPIN 300MG CAPSULES**  
QTY: 60    2 REFILLS BEFORE 11/29/11  
Refill      NDC:00527-1315-30  
Retail Price: \$113.89    Your Insurance Saved You: \$108.89    \$ 5.00

A: JAIN, MD      PLAN: PAIDMPD  
MFG: LANNETT      GROUP: CMD3896  
XXX/KHN/KHN/KHN/RJW      CLAIM REF# ONA1TWP

**Walgreens**      12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

SUN 12:00PM  
Refill

RIFAMPIN 300MG CAPSULES  
00527-1315-30  
ALPHA

QTY 60  
20 DRAM



**DARK REDDISH-BROWN**  
**FRONT: LANNETT**  
**BACK: 1315**

XXX/KHN/KHN/KHN/RJW

Brunsting004477



## Are you getting the most from your Medicare Part D plan?

We can help you find out with a **FREE one-on-one plan review session** with your pharmacist. We'll review your medications, look for ways to help save you money and more.

*Make your appointment today!*

**Walgreens**


There's a way to stay well.

Information provided by Walgreens is not a substitute for your own research concerning your healthcare, including Medicare.

04589

## Prescription Transfers Easy as 1-2-3

1. Gather your prescription bottles.
2. Call or visit your local Walgreens.
3. We'll take care of the rest!



Avoid potentially harmful drug interactions by filling all your prescriptions at one pharmacy. By choosing Walgreens, our pharmacists can screen your prescriptions and warn you of potential interactions.

**For the Walgreens nearest you,  
call 1-800-WALGREENS  
or visit [Walgreens.com/findastore](http://Walgreens.com/findastore).**

Another great service from

**Walgreens**  
The Pharmacy • America Trusts • Since 1901®

The Pharmacy • America Trusts • Since 1901®

OPT-2875 677 002 024080

INPO-0990 00527131530

38680

8  
BR



NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

- Store TT 4 of 5
- Your Insurance Saved You: \$108.89

SUN 12:00PM  
\$5.00  
EXPRESS PAY

01/01/11  
Refill



## Personal Prescription Information

**LOOK INSIDE FOR IMPORTANT INFORMATION**

**ABOUT YOUR MEDICATION.**

Take advantage of this convenient service:

• **Auto Refills**

We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

**Walgreens**

The Pharmacy • America Trusts • Since 1901®

Visit us online at [Walgreens.com](http://Walgreens.com)

***Thank you for choosing Walgreens!***

Brunsting004478

# YOUR PERSONAL PRESCRIPTION INFORMATION

**Your Walgreens Pharmacy Location**  
 12850 Memorial Drive  
 Houston, TX 77024  
 (713)722-7247

<b>PATIENT</b> NELVA BRUNSTING	<b>DOCTOR</b> A. JAIN, MD	<b>DRUG DESCRIPTION</b>
<b>BIRTH DATE</b> 10/08/26		
<b>MEDICATION</b> SPIRIVA CAPS 30'S & HANDIHALER	<b>PATIENT ALLERGIES</b>	
<b>QUANTITY</b> 30		
<b>DIRECTIONS</b> INHALE CONTENTS OF ONE CAPSULE ONCE DAILY USING HANDIHALER		

**INGREDIENT NAME:** TIOTROPIUM BROMIDE (TYE-oh-TROE-pee-um BROE-mide)

**COMMON USES:** This medicine is an anticholinergic agent used to treat the symptoms of chronic obstructive pulmonary disease, including chronic bronchitis and emphysema. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking another anticholinergic (eg, ipratropium). Ask your doctor if you are unsure if any of your medicines are anticholinergics. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medicine you are taking if you have trouble urinating; an enlarged prostate; bladder blockage; glaucoma; kidney problems; allergies (including milk proteins); pregnancy; or breast-feeding. Tell your doctor if you are having an asthma attack or increased difficulty breathing. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. An additional patient information leaflet is available with this medicine. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have before using this medicine. **DO NOT SWALLOW THE CAPSULES. THE CAPSULES ARE USED WITH A SPECIAL INHALER.** Do NOT remove a capsule from the packaging until you are ready to use it. To remove a capsule from the packaging, carefully peel the foil back to expose the capsule, then tip the capsule out of the blister. Do NOT cut the foil or use sharp objects to remove the capsule from the blister. If a second capsule is exposed to the air when you are removing a capsule for use, it must be discarded. Do not save the capsule for later. Place the capsule in the special inhaler device immediately. The device will puncture the capsule so that the medicine inside may be inhaled into the lungs through the mouthpiece. Exhale slowly and deeply. Do NOT breathe into the mouthpiece of the inhaler. Position the inhaler mouthpiece between your lips and try to rest your tongue flat. Keep your head upright. Do not block the air vents on the inhaler. **TAKE A SLOW, DEEP BREATH.** You should hear or feel the capsule vibrate inside the inhaler. Hold your breath as long as it feels comfortable, then exhale slowly through pursed lips. Breathe out completely. To be sure all of the medicine has been inhaled from the capsule, inhale from the mouthpiece a second time following the same process. **ASK YOUR DOCTOR OR PHARMACIST** if you are unclear on how to use this device or inhale the medicine. **CLEAN THE INHALER DEVICE** once a month according to the instructions in the patient leaflet. Each inhaler device may be used for up to 1 year and then should be replaced. **STORE THIS MEDICINE** at room temperature at 77 degrees F (25 degrees C), away from moisture and light. Brief storage between 59 and 86 degrees F (15 and 30 degrees C) is permitted. Do not store in the bathroom. This medicine will work best if used regularly. Taking your dose at the same time each day will help you to remember. **IF YOU MISS A DOSE** OF THIS MEDICINE, use it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do NOT use 2 doses at once. Do NOT use this medicine more often than 1 time every 24 hours.

**CAUTIONS:** DO NOT USE THIS MEDICINE if you are allergic to any ingredient in this medicine or to atropine or related medicines (eg, ipratropium). **THIS MEDICINE WILL NOT STOP AN ASTHMA ATTACK** once one has started. **IF YOU ARE ALSO USING A RESCUE INHALER** (eg, albuterol), be sure to always carry the bronchodilator inhaler with you to use during asthma attacks. **DO NOT EXCEED THE RECOMMENDED DOSE** without checking with your doctor. **DO NOT STOP USING THIS MEDICINE** without first checking with your doctor. This medicine may cause dizziness or blurred vision. These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. **AVOID GETTING THIS MEDICINE IN YOUR EYES.** If you get the medicine in your eyes and eye pain, blurred vision, or other vision changes occur, contact your doctor immediately. **THIS MEDICINE MAY SOMETIMES CAUSE SEVERE BREATHING PROBLEMS** right after you use a dose. If this happens, use your short-acting bronchodilator inhaler. Contact your doctor or seek other medical care at once. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE,** either prescription or over-the-counter, check with your doctor or pharmacist. **USE THIS MEDICINE WITH CAUTION IN THE ELDERLY;** they may be more sensitive to its effects, especially constipation and urinary tract infections. **FOR WOMEN: IF YOU**

**PLAN ON BECOMING PREGNANT,** discuss with your doctor the benefits and risks of using this medicine during pregnancy. **IT IS NOT KNOWN IF THIS MEDICINE IS FOUND** in breast milk. **IF YOU ARE OR WILL BE BREAST-FEEDING** while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while using this medicine include dry mouth, constipation, sinus inflammation, throat irritation, stomach pain, vomiting, blurred vision, or mild nosebleed. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience irritation, pain, or white patches in your mouth or on your tongue; mouth sores; severe or persistent nosebleeds; severe or persistent constipation; difficult or painful urination; eye pain or discomfort; vision changes (eg, halos, colored images); chest pain; fast or irregular heartbeat; difficulty swallowing; new or worsened breathing problems; or wheezing. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include mental changes; severe constipation; stomach pain; or tremors.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. **KEEP THIS MEDICINE** FOR the reach of children and pets. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** obtain refills before your supply runs out.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

**KEEP OUT OF REACH OF CHILDREN. STORE IN SAFETY CONTAINER OR SECURE AREA.**

**NELVA BRUNSTING**  
 13630 Pinrock, Houston, TX 770797517  
 (713)464-4391

**RX # 1496586-03328** DATE: 01/01/11

**SPIRIVA CAPS 30'S & HANDIHALER**  
 QTY: 30 2 REFILLS BEFORE 12/05/11  
 Refill NDC:00597-0075-41  
 Retail Price: \$260.99 Your Insurance Saved You: \$44.45

\$ 216.54

A. JAIN, MD PLAN: PAID/PPD  
 MFG:BOEHRINGER GROUP# CND3898  
 XXX/JIC/JIC/ /RJW CLAIM REF# MRWHDCC

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
 PH: (713)722-7247

Customer Receipt

**NELVA BRUNSTING**  
 13630 Pinrock, Houston, TX 770797517  
 (713)464-4391

**RX # 1496586-03328** DATE: 01/01/11

**SPIRIVA CAPS 30'S & HANDIHALER**  
 QTY: 30 2 REFILLS BEFORE 12/05/11  
 Refill NDC:00597-0075-41  
 Retail Price: \$260.99 Your Insurance Saved You: \$44.45

\$ 216.54

A. JAIN, MD PLAN: PAID/PPD  
 MFG:BOEHRINGER GROUP# CND3898  
 XXX/JIC/JIC/ /RJW CLAIM REF# MRWHDCC

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
 PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

SUN 12:00PM  
 Refill

SPIRIVA CAPS 30'S & HANDIHALER  
 QTY 30  
 00597-0075-41  
 ALPHA

XXX/JIC/JIC/ /RJW

Brunsting004469

AABrunsting.Financials003952

Get a **FREE** one-on-one  
**Medicare Part D**  
review session!

Your pharmacist will:

- Review your medications and look for ways to help save you money
- Talk about other things you can do to stay well
- Check for potential drug interactions



Make your appointment today!

**Walgreens**

There's a way to stay well.

Information provided by Walgreens is not a substitute for your own research concerning your healthcare, including Medicare.

02739

**Don't take  
chances with  
your health.**

Fill all your prescriptions  
at one pharmacy

Walgreens automatically screens for drug interactions, including any over-the-counter medicines or supplements you tell us about.



**Play it safe. Fill all your  
prescriptions at Walgreens.**

OFF: 28752677 002 0240098

INFO: 0990 00397007541

3021

8  
BR




NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

- Store TT 5 of 5
- Your Insurance Saved You: \$44.45

SUN 12:00PM  
**\$216.54**  
EXPRESS PAY

01/01/11  
Refill



## Personal Prescription Information

**LOOK INSIDE FOR IMPORTANT INFORMATION**

**ABOUT YOUR MEDICATION.**

Take advantage of this convenient service:

- **Auto Refills**  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

**Walgreens**

The Pharmacy America Trusts • Since 1901™

Visit us online at [Walgreens.com](http://Walgreens.com)

**Thank you for choosing Walgreens!**

Brunsting004470

# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
12850 Memorial Drive  
Houston, TX 77024  
(713)722-7247

<b>PATIENT</b>	NELVA BRUNSTING	<b>DOCTOR</b>	A. JAIN, MD	<b>DRUG DESCRIPTION</b>	
<b>BIRTH DATE</b>	10/08/26	<b>PATIENT ALLERGIES</b>		<b>LIQUID</b>	
<b>MEDICATION</b>	BROVANA 15MCG/2ML INH SOL 30X2ML				
<b>QUANTITY</b>	120				
<b>DIRECTIONS</b>	INHALE 1 VIAL VIA NEBULIZER TWOX DAILY				

**INGREDIENT NAME:** ARFORMOTEROL (ar-for-MOE-ter-ol)

**COMMON USES:** This medicine is a long-acting beta-agonist bronchodilator used for long-term treatment of chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** **WARNING: LONG-ACTING BETA-AGONISTS SUCH AS THIS MEDICINE HAVE BEEN RARELY ASSOCIATED WITH AN INCREASED RISK OF ASTHMA-RELATED DEATH.** Long-acting beta-agonists should not be used in asthma patients without another long-term asthma-control medicine (eg, inhaled corticosteroids). This medicine has not been approved to treat asthma. Safety and effectiveness of this medicine in asthma patients have not been confirmed. **SOME MEDICINES MAY INTERACT** with this medicine. **DO NOT TAKE THIS MEDICINE** if you are using another medicine that has a long-acting beta-agonist (eg, salmeterol) in it. **TELL YOUR HEALTH CARE PROVIDER** if you are taking any other medicine, especially any of the following: corticosteroids (eg, prednisone), diuretics (eg, furosemide, hydrochlorothiazide), xanthines (eg, theophylline), catechol-O-methyltransferase (COMT) inhibitors (eg, entacapone), monoamine oxidase inhibitors (MAOI) (eg, phenelzine), or beta-blockers (eg, propranolol). This may not be a complete list of all interactions that may occur. Tell your doctor if you have taken MAOI (eg, phenelzine) or a tricyclic antidepressant (eg, amitriptyline) within the last 14 days. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you are having severe breathing problems (eg, sudden, severe onset or worsening of COPD symptoms such as chest tightness, cough, shortness of breath, wheezing), you have asthma and you are not currently using a long-term asthma-control medicine (eg, inhaled corticosteroids), or if you have asthma that is already well controlled with the use of a long-term asthma-control medicine. **THIS MEDICINE SHOULD NOT BE USED IN CHILDREN;** safety and effectiveness in children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine comes with a MEDICATION GUIDE approved by the U.S. Food and Drug Administration. Read it carefully each time you refill this medicine. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. **THIS MEDICINE SHOULD ONLY BE INHALED USING A NEBULIZER.** Do NOT inject or swallow it. A health care provider will teach you how to use the nebulizer. Be sure you know what type of nebulizer to use with this medicine, and how to use it. Follow the procedure you are taught when you use a dose. Contact your health care provider if you have any questions. **STORE THIS MEDICINE** in the original foil pouch in a dry place. Do not remove from the foil pouch of the vial until right before use. **IF THE MEDICINE CONTAINS PARTICLES,** is cloudy or discolored, or if the vial is cracked or damaged in any way, do not use it. **DO NOT MIX THIS MEDICINE WITH OTHER MEDICINES** in your nebulizer machine. **TO USE THIS MEDICINE,** twist open the top of the vial and pour the entire contents into the nebulizer reservoir. **CONNECT THE NEBULIZER RESERVOIR** to the mouthpiece or face mask. **CONNECT THE NEBULIZER** to the compressor. Sit in a comfortable, upright position. **PLACE THE MOUTHPIECE** in your mouth (or put on the face mask) and turn on the nebulizer. **BREATHE AS CALMLY,** deeply, and evenly as possible. **INHALE** as formed in the nebulizer chamber (about 5 to 10 minutes). **CLEAN THE NEBULIZER** according to the instructions. Failure to properly clean the nebulizer could lead to bacteria entering the medicine. This may lead to an infection. To avoid bacteria entering the medicine, the entire contents of the nebulizer should be used on the first time. **STORE THIS MEDICINE IN THE REFRIGERATOR,** between 36 and 46 degrees F (2 and 8 degrees C). Do not freeze. **UNOPENED POUCHES MAY BE STORED** at room temperature between 68 and 77 degrees F (20 and 25 degrees C) for up to 6 weeks. Store away from heat, moisture, and light. If this medicine is stored from room temperature for up to 6 weeks, **DO NOT USE THIS MEDICINE** if it is past the expiration date on the container. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **CONTINUE TO USE THIS MEDICINE** even if you feel well. Do not miss any doses. If you miss a dose of this medicine, skip the missed dose and go back to your regular dosing schedule. Do not use 2 doses at once.

**CAUTIONS:** **DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine or to formoterol. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescription or over the counter, check with your doctor or pharmacist. **LAB TESTS,** including lung function and blood potassium levels, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to keep all doctor and lab appointments. **THIS MEDICINE MAY CAUSE DIZZINESS.** This effect may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. Do not drive or perform other, possibly unsafe tasks until you know how you react to it. **THIS MEDICINE WILL NOT STOP SUDDEN SYMPTOMS OF COPD ONCE THEY HAVE ALREADY STARTED.** Be sure to carry a short-acting bronchodilator inhaler (eg, albuterol) with you at all times to treat any breathing problems that may occur between doses of this medicine (eg, severe or sudden onset of wheezing or shortness of breath). If you have any questions about which medicines stop sudden symptoms, check with your doctor or pharmacist. **IF YOU HAVE BEEN REGULARLY USING A SHORT-ACTING BRONCHODILATOR INHALER,** you may experience some changes about how to use it with this medicine. Short-acting bronchodilators are normally used with this medicine to treat breathing problems that may occur between doses. **THE RISK OF SERIOUS HEART PROBLEMS** (eg, irregular heartbeat) may be greater if you use this medicine in high doses. Do NOT use more than recommended dose or use more often than prescribed. **TELL YOUR DOCTOR AT ONCE** if you notice that your short-acting bronchodilator inhaler does not work as well, if you need to use it often, or if your breathing problems get worse. **CONTACT YOUR DOCTOR OR SEEK MEDICAL CARE RIGHT AWAY** if you have breathing problems that worsen quickly, or if you use your short-acting bronchodilator and do not get relief. **TALK WITH YOUR DOCTOR OR PHARMACIST** about all of your breathing medicines and how to use them. Do not start, stop, or change the dose of any

breathing medicine unless your doctor tells you to. **THE MEDICINE MAY SOMETIMES CAUSE SEVERE BREATHING PROBLEMS RIGHT AFTER YOU USE A DOSE.** If this happens, use your short-acting bronchodilator. **CONTACT YOUR DOCTOR OR SEEK OTHER MEDICAL CARE AT ONCE.** **THIS MEDICINE MAY RAISE YOUR BLOOD SUGAR.** High blood sugar may make you feel confused, drowsy, or thirsty. It can also make you flush, breathe faster, or have a fruit-like breath odor. If these symptoms occur, tell your doctor right away. **PREGNANCY and BREAST-FEEDING:** If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. It is not known if this medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby.

**POSSIBLE SIDE EFFECTS:** **SIDE EFFECTS** that may occur while taking this medicine include: back pain; diarrhea; dry mouth; headache; nausea; nervousness; stuffy nose; tiredness; tremor; trouble sleeping; vomiting. If they continue or are bothersome, check with your doctor. **SEEK MEDICAL ATTENTION RIGHT AWAY** if any of these SEVERE side effects occur: chest pain; fast or irregular heartbeat; fever, chills, or persistent sore throat; leg swelling; new or worsening breathing problems (eg, increased chest tightness, coughing, shortness of breath, wheezing); severe or persistent headache, dizziness, tremor, or nervousness; severe or persistent muscle pain or cramps; symptoms of high blood sugar (eg, increased thirst, urination, or hunger; unusual weakness or drowsiness; confusion); trouble speaking. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, tongue, or throat; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** **IF OVERDOSE IS SUSPECTED,** contact your local poison control center or emergency room immediately. Symptoms may include chest pain; fast or irregular heartbeat; severe or persistent dizziness; dry mouth; fatigue; headache; muscle pain or cramps; nausea; nervousness; trouble sleeping, or tremor; severe or persistent symptoms of high blood sugar (eg, increased thirst, urination, or hunger; drowsiness; flushing of the skin; confusion; fruit-like breath odor).

**ADDITIONAL INFORMATION:** If your symptoms do not improve or if they become worse, check with your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **USING THIS MEDICINE FOR AN EXTENDED PERIOD** of time may obtain refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 959810

KEEP OUT OF REACH OF CHILDREN. STORE IN SAFETY CONTAINER OR SECURE AREA.

**NELVA BRUNSTING**

12630 Pinecroft, Houston, TX 770797517  
(713)464-4321

**RX # 1496587-03328**

DATE: 01/13/11

**BROVANA 15MCG/2ML INH SOL 30X2ML**  
QTY: 120 2 REFILLS BEFORE 12/05/11  
Refill NDC: 63402-0911-30

\$ 527.99

A. JAIN, MD  
MFG:SEPRACOR  
JDC/SSH/SSH/IKSC

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Customer Receipt

**NELVA BRUNSTING**

12630 Pinecroft, Houston, TX 770797517  
(713)464-4321

**RX # 1496587-03328**

DATE: 01/13/11

**BROVANA 15MCG/2ML INH SOL 30X2ML**  
QTY: 120 2 REFILLS BEFORE 12/05/11  
Refill NDC: 63402-0911-30

\$ 527.99

A. JAIN, MD  
MFG:SEPRACOR  
JDC/SSH/SSH/IKSC

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

THU 4:15PM  
Refill

BROVANA 15MCG/2ML INH SOL 30X2ML  
63402-0911-30  
REFRIG

QTY 120

LIQUID

JDC/SSH/SSH/IKSC

Med Guide

Brnusting004471

AABrunsting.Financials003954


Don't take a chance with your lungs — get a PARI Nebulizer cup now.



Did you know a PARI nebulizer was used to prove safety and efficacy of your inhaled medication? PARI nebulizers are the clinical choice, make them your choice.


**The Benefits of PARI Reusable Nebulizers**

- Short treatment times
- Consistent, efficient delivery of medication to the lungs
- Reusable, designed to last 6 months
- Easy to clean: boilable and dishwasher safe
- Cost effective



PARI LCP Plus used in clinical trial.

**Continue to effectively manage your COPD by using a PARI Nebulizer.**



Chronic obstructive pulmonary disease (COPD) is with you all of the time, even when you don't have symptoms. That is why it's important to use the PARI Nebulizer system, to deliver your medication to you effectively.

**If you're not using a PARI nebulizer cup, you may not be getting the most benefit from your aerosol medication.**

PARI reusable nebulizers feature breath enhanced technology resulting in increased aerosol delivery with decreased medication waste.

For more information contact PARI at 1.800.FAST.NEB (327.8632) or your Walgreens pharmacist - visit [www.PARI.com](http://www.PARI.com).

**Talk to your doctor or pharmacist today about the PARI Reusable Nebulizer cup.**

This message was developed and paid for by PARI Respiratory Equipment, Inc.

8  
BR




\*1496587 0902 3 0062799 0\*

THU 4:15PM  
\$527.99  
EXPRESS PAY

NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

01/13/11  
Refill  
REFRIGERATE

- MEDICARE: Patient Not Eligible- Use New Plan or charge Cash
  - Store TT 1 of 1
  - Prescription Savings Club could save you \$117.29!
- Ask if you qualify.



**Personal Prescription Information**

**LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.**

**Take advantage of these convenient services:**

- **Touch Tone Refills**  
Save time by using our automated system for a refill. Just dial the number on your prescription label.
- **Auto Refills**  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.



Visit us online at [Walgreens.com](http://Walgreens.com)

***Thank you for choosing Walgreens!***

To stop receiving sponsored info at this pharmacy, call 888-336-5744 or visit [remove-me.net](http://remove-me.net) Use code:2875677 013 0320825

Brunsting004472



# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
12850 Memorial Drive  
Houston, TX 77024  
(713)722-7247

<b>PATIENT</b>	NELVA BRUNSTING	<b>DOCTOR</b>	A. JAIN, MD	<b>DRUG DESCRIPTION</b>	
<b>BIRTH DATE</b>	10/08/26	<b>PATIENT ALLERGIES</b>			LIQUID
<b>MEDICATION</b>	BROVANA 15MCG/2ML INH SOL 30X2ML				
<b>QUANTITY</b>	120				
<b>DIRECTIONS</b>	INHALE 1 VIAL VIA NEBULIZER TWICE DAILY				

**INGREDIENT NAME:** ARFORMOTEROL (ar-for-MOE-ter-ole)

**COMMON USES:** This medicine is a long-acting beta-agonist bronchodilator used for long-term treatment of chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** **WARNING: LONG-ACTING BETA-AGONISTS SUCH AS THIS MEDICINE HAVE BEEN BARELY ASSOCIATED WITH AN INCREASED RISK OF ASTHMA-RELATED DEATH.** Long-acting beta-agonists should not be used in asthma patients not on asthma-control medicine (eg, inhaled corticosteroids). This medicine has not been approved to treat asthma. Safety and effectiveness of this medicine in patients with asthma have not been confirmed. **SOME MEDICINES MAY INTERACT** with this medicine. **DO NOT TAKE THIS MEDICINE** if you are using another medicine that has a long-acting beta-agonist (eg, salmeterol) in it. **TELL YOUR HEALTH CARE PROVIDER** if you are taking any other medicines, especially any of the following: corticosteroids (eg, prednisone), medicine (eg, tricyclic antidepressants, xanthines (eg, theophylline), catechol-O-methyltransferase (COMT) inhibitors (eg, entacapone), monoamine oxidase inhibitors (MAOIs) (eg, phenelzine, tricyclic antidepressants (eg, amitriptyline), or beta-blockers (eg, propranolol). **DO NOT STOP OR STOP ANY MEDICINE** without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including a history of other breathing problems (eg, asthma), diabetes, heart problems (eg, fast or irregular heartbeat, heart blood vessel problems), liver problems, high blood pressure, low blood potassium levels, seizures, an overactive thyroid, or you have high blood or urine ketone levels, allergies, pregnancy, or breast-feeding. Tell your doctor if you have recently been to an emergency room for breathing problems; have a history of frequent hospitalizations for breathing problems; have ever had life-threatening breathing problems; or have had an unusual reaction to a sympathomimetic medicine (eg, albuterol, pseudoephedrine), such as fast or irregular heartbeat, overexcitement, or severe trouble sleeping. Tell your doctor if you have taken a MAOI (eg, phenelzine) or a tricyclic antidepressant (eg, amitriptyline) within the last 14 days. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you are having severe breathing problems (eg, sudden, severe onset or worsening of COPD symptoms such as chest tightness, cough) or if you have asthma that is already well controlled with the use of a long-term asthma-control medicine. **THIS MEDICINE SHOULD NOT BE USED IN CHILDREN** safety and effectiveness in children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine comes with a MEDICATION GUIDE approved by the U.S. Food and Drug Administration. Read it carefully each time you refill this medicine. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. **THIS MEDICINE SHOULD ONLY BE INHALED USING A NEBULIZER.** Do NOT inject or swallow it. A health care provider will teach you how to use the nebulizer. Be sure you know what type of nebulizer to use with this medicine, and how to use it. Follow the procedures you are taught when you use a dose. Contact your health care provider if you have any questions. **STOP USING THIS MEDICINE** in the original foil pouch in a dry place. Do not remove the foil pouch or the vial until right before use. **IF THE MEDICINE CONTAINS PARTICLES** is cloudy or discolored, or if the vial is cracked or damaged in any way, do not use it. **DO NOT MIX THIS MEDICINE WITH OTHER MEDICINES** in your nebulizer machine. **DO NOT USE THIS MEDICINE** until you see the top of the vial and pour the entire contents into the nebulizer reservoir. **CONNECT THE NEBULIZER RESERVOIR** to the mouthpiece or face mask. **CONNECT THE MOUTHPIECE** to the compressor. Sit in a comfortable, upright position. **PLACE THE MOUTHPIECE** in your mouth (or put on the face mask) and turn on the nebulizer. **BREATHE IN CALMLY,** deeply, and evenly as possible until no more mist is formed in the nebulizer chamber (about 5 to 10 minutes). **CLEAN THE NEBULIZER** according to the instructions. Failure to properly clean the nebulizer could lead to bacteria entering the medicine. This may lead to an infection. To avoid bacteria entering the medicine, use the entire contents right after opening the vial for the first time. **DO NOT STORE THIS MEDICINE** in the refrigerator, between 36 and 46 degrees F (2 and 8 degrees C). Do not freeze. **UNOPENED BOTTLES MAY BE STORED** at room temperature between 59 and 77 degrees F (10 and 25 degrees C) for up to 6 weeks. Store away from heat, moisture, and light. If this medicine is stored at room temperature, throw it away after 6 weeks. **DO NOT USE** this medicine if it is past the expiration date on the container. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **CONTINUE TO USE THIS MEDICINE** even if you feel well. Do not miss any doses. If you miss a dose of this medicine, skip the missed dose and go back to your regular dosing schedule. Do not use 2 doses at once.

**CAUTIONS:** **DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine or to formoterol. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescription or over-the-counter, check with your doctor or pharmacist. **LAB TESTS,** including lung function and blood potassium levels, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to keep all doctor and lab appointments. **THIS MEDICINE MAY CAUSE DIZZINESS.** This effect may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. Do not drive or perform other possibly unsafe tasks until you know how you react to it. **THIS MEDICINE DOES NOT STOP ASTHMA OR COPD. IF YOU ARE ALREADY TAKING THEM, HAVE ALREADY STARTED,** Be sure to carry a short-acting bronchodilator inhaler (eg, albuterol) with you at all times to treat any breathing problems that may occur between doses of this medicine (eg, severe or sudden onset of wheezing or shortness of breath). If you have any questions about which medicines you should use, check with your doctor or pharmacist. **IF YOU HAVE BEEN IRREGULARLY USING A SHORT-ACTING BRONCHODILATOR INHALER,** talk with your doctor about how to use it with this medicine. Short-acting bronchodilators are normally used with this medicine to treat breathing problems that may occur between doses. **THE RISK OF SERIOUS HEART PROBLEMS IS GREATER** if you use this medicine in high doses. Do NOT use more than recommended dose or use more often than prescribed. **TELL YOUR DOCTOR AT ONCE** if you notice that your short-acting bronchodilator inhaler does not work as well, if you need to use it often, or if your breathing problems get worse. **CONTACT YOUR DOCTOR FOR OR SEEK MEDICAL CARE RIGHT AWAY** if you have breathing problems that worsen quickly, or if you use your short-acting bronchodilator and do not get relief. **TALK WITH YOUR DOCTOR OR PHARMACIST** about all of your breathing medicines and how to use them. Do not start, stop, or change the dose of any

breathing medicine unless your doctor tells you to. **THE MEDICINE MAY, SOMETIMES CAUSE SEVERE BREATHING PROBLEMS RIGHT AFTER YOU USE A DOSE.** If this happens, use your short-acting bronchodilator. Contact your doctor or seek other medical care at once. **THIS MEDICINE MAY RAISE YOUR BLOOD SUGAR.** High blood sugar may make you feel confused, drowsy, or thirsty. It can also make you flush, breathe faster, or have a fruit-like breath odor. If these symptoms occur, tell your doctor right away. **PREGNANCY AND BREAST-FEEDING:** If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. It is not known if this medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby.

**POSSIBLE SIDE EFFECTS:** **SIDE EFFECTS** that may occur while taking this medicine include: back pain; diarrhea; dry mouth; headache; nausea; nervousness; stuffy nose; tiredness; tremor; trouble sleeping; vomiting. If they continue or are bothersome, check with your doctor. **SEEK MEDICAL ATTENTION RIGHT AWAY** if any of these **SEVERE** side effects occur: chest pain; fast or irregular heartbeat; fever, chills, or persistent sore throat; leg swelling; new or worsening breathing problems (eg, increased chest tightness, coughing, shortness of breath, wheezing); severe or persistent headache, dizziness, tremor, or nervousness; severe or persistent muscle pain or cramps; symptoms of high blood sugar (eg, increased thirst, urination, or hunger; unusual weakness or drowsiness; confusion); trouble speaking. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, tongue, or throat; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** **IF OVERDOSE IS SUSPECTED,** contact your local poison control center or emergency room immediately. Symptoms may include chest pain; fast or irregular heartbeat; severe or persistent dizziness, dry mouth, fatigue, headache, muscle pain or cramps, nausea, nervousness, trouble sleeping, or tremor; severe or persistent symptoms of high blood sugar (eg, increased thirst, urination, or hunger; drowsiness; flushing of the skin; confusion; fruit-like breath odor).

**ADDITIONAL INFORMATION:** If your symptoms do not improve or if they become worse, check with your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** obtain refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

**NELVA BRUNSTING**  
13630 Pinerock, Houston, TX 770797517  
(713)464-4391  
**RX # 1496587-03328** DATE: 02/18/11

**BROVANA 15MCG/2ML INH SOL 30X2ML**  
QTY: 120 1 REFILL BEFORE 12/05/11  
Refill NDC:63402-0911-30 \$ 527.99

A. JAIN, MD  
MFG:SEPRACOR  
HMC / / / KDM

**Walgreens** 12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Customer Receipt

**NELVA BRUNSTING**  
13630 Pinerock, Houston, TX 770797517  
(713)464-4391  
**RX # 1496587-03328** DATE: 02/18/11

**BROVANA 15MCG/2ML INH SOL 30X2ML**  
QTY: 120 1 REFILL BEFORE 12/05/11  
Refill NDC:63402-0911-30 \$ 527.99

A. JAIN, MD  
MFG:SEPRACOR  
HMC / / / KDM

**Walgreens** 12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

BROVANA 15MCG/2ML INH SOL 30X2ML	QTY 120	LIQUID
TUE 12:00PM	63402-0911-30	
Refill	REFRIG	

HMC / / / KDM Med Guide

Brnstin004467


Don't take a chance with your lungs — get a PARI Nebulizer cup now.



Did you know a PARI nebulizer was used to prove safety and efficacy of your inhaled medication? PARI nebulizers are the clinical choice, make them your choice.


**The Benefits of PARI Reusable Nebulizers**

- Short treatment times
- Consistent, efficient delivery of medication to the lungs
- Reusable, designed to last 6 months
- Easy to clean: boilable and dishwasher safe
- Cost effective



PARI LC® Plus used in clinical trial.

3544



Continue to effectively manage your COPD by using a PARI Nebulizer.

Chronic obstructive pulmonary disease (COPD) is with you all of the time, even when you don't have symptoms. That is why it's important to use the PARI Nebulizer system, to deliver your medication to you effectively.

If you're not using a PARI nebulizer cup, you may not be getting the most benefit from your aerosol medication.

PARI reusable nebulizers feature breath enhanced technology resulting in increased aerosol delivery with decreased medication waste.



Talk to your doctor or pharmacist today about the PARI Reusable Nebulizer cup.

This message was developed and paid for by PARI Respiratory Equipment, Inc.

OCT-2875 677 049 0537281

INFO: 0806.63402091130

3544

8 BR



\*1486587 2203 3 0062798 0\*

NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

- MEDICARE: Patient Not Eligible- Use New Plan or charge Cash
  - Store TT 1 of 1
  - Prescription Savings Club could save you \$117.29!
- Ask if you qualify.

TUE 12:00PM  
**\$527.99**  
EXPRESS PAY

02/18/11

Refill

REFRIGERATE  
MED GUIDE



Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.

Take advantage of these convenient services:

- **Auto Refills**  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.
- **Touch Tone Refills**  
Save time by using our automated system for a refill. Just dial the number on your prescription label.

*Walgreens*

The Pharmacy America Trusts • Since 1901™

Visit us online at Walgreens.com


*Thank you for choosing Walgreens!*

To stop receiving sponsored info at this pharmacy, call 888-336-5744 or visit remove-me.net Use code:2875677 049 0537281

Brunsting004468

# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
12850 Memorial Drive  
Houston, TX 77024  
(713)722-7247

<b>PATIENT</b> NELVA BRUNSTING	<b>DOCTOR</b> A. JAIN, MD	<b>DRUG DESCRIPTION</b>
<b>BIRTH DATE</b> 10/08/26		
<b>MEDICATION</b> AZITHROMYCIN 250MG TABLETS	<b>PATIENT ALLERGIES</b>	
<b>QUANTITY</b> 30		<b>PINK</b>
<b>DIRECTIONS</b> TAKE 1 TABLET BY MOUTH EVERY DAY		<b>FRONT: 93</b>
		<b>BACK: 7146</b>

**INGREDIENT NAME:** AZITHROMYCIN  
(ay-ZITH-roe-MYE-sin)

**COMMON USES:** This medicine is a macrolide antibiotic used to treat bacterial infections.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking dofetilide, nilotinib, propafenone, pirozide, or tetrabenazine. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking antiarrhythmics (eg, disopyramide), anticoagulants (eg, warfarin), arsenic, astemizole, carbamazepine, cisapride, digoxin, domperidone, meprobamate, methadone, nelfinavir, cyclosporine, ergot alkaloids (eg, ergotamine), paliperidone, phenytoin, quinolones (eg, levofloxacin), rifampin, terfenadine, theophylline, triazolam, tyrosine kinase inhibitors (eg, dasatinib), or medicines that may affect your heartbeat. Ask your doctor if you are unsure if any of the medicines you are taking may affect your heartbeat. Inform your doctor of any other medical conditions including irregular heartbeat, kidney problems, liver problems, myasthenia gravis, allergies, pregnancy or breastfeeding. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine may be taken on an empty stomach or with food. DO NOT TAKE THIS MEDICINE within 1 hour before or 2 hours after aluminum- or magnesium-containing antacids. STORE THIS MEDICINE at room temperature, away from heat and light. TO CLEAR UP YOUR INFECTION COMPLETELY, continue taking this medicine for the full course of treatment even if you feel better in a few days. Do not miss any doses. Taking this medicine at the same time each day will make it easier to remember. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. If you miss a dose, do not take 2 doses at once.

**CAUTIONS:** DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it, to other macrolide antibiotics (such as erythromycin), to ketolide antibiotics (such as telithromycin), or if you are allergic to any ingredient in this product. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to erythromycin or any macrolide or ketolide antibiotic. A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine, contact your doctor or pharmacist. IF YOU EXPERIENCE difficulty breathing; tightness of chest; swelling of eyelids, face, or lips; or if you develop a rash or hives, tell your doctor immediately. Do not take any more of this medicine unless your doctor tells you to do so. This medicine may cause drowsiness, dizziness, or lightheadedness. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. THIS

MEDICINE MAY CAUSE increased sensitivity to the sun. Avoid exposure to the sun, sunlamps, or tanning booths until you know how you react to this medicine. Use a sunscreen or protective clothing if you must be outside for a prolonged period. MILD DIARRHEA IS COMMON with antibiotic use. However, a more serious form of diarrhea (pseudomembranous colitis) may rarely occur. This may develop while you use the antibiotic or within several months after you stop using it. Contact your doctor right away if stomach pain or cramps, severe diarrhea, or bloody stools occur. Do not treat diarrhea without first checking with your doctor. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. FOR WOMEN: IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. IT IS UNKNOWN IF THIS MEDICINE IS EXCRETED in breast milk. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS, that may go away during treatment, include mild diarrhea, headache, nausea, or stomach pain or upset. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience vomiting, changes in hearing, or ringing in the ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience chest pain; swelling of your hands, legs, face, lips, eyes, throat, or tongue; difficulty swallowing, speaking, or breathing; eye or vision problems; fainting; hearing loss; irregular heartbeat; muscle weakness; reddened, blistered, or swollen skin; seizures; severe stomach pain, cramps, or diarrhea; bloody stools; unusual vaginal itching, odor, or discharge; or yellowing of the eyes or skin. An allergic reaction to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, dizziness, trouble breathing, or unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include nausea, vomiting, stomach upset, and diarrhea.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

NELVA BRUNSTING  
13830 Pinerock, Houston, TX 770797517  
(713)464-4391  
RX # 1494789-03328 DATE: 03/02/11

AZITHROMYCIN 250MG TABLETS  
QTY: 30 NO REFILLS - DR. AUTH REQUIRED  
Refill NDC:00093-7146-56  
Retail Price: \$195.79 Your Insurance Saved You: \$190.79

\$ 5.00

A. JAIN, MD MFG:TEVA PLAN: PAIDMPD  
XXXI / KMN/KMN GROUP# CMD3896  
CLAIM REF# FKRWQHL

**Walgreens**  
12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Customer Receipt

NELVA BRUNSTING  
13830 Pinerock, Houston, TX 770797517  
(713)464-4391  
RX # 1494789-03328 DATE: 03/02/11

AZITHROMYCIN 250MG TABLETS  
QTY: 30 NO REFILLS - DR. AUTH REQUIRED  
Refill NDC:00093-7146-56  
Retail Price: \$195.79 Your Insurance Saved You: \$190.79

\$ 5.00

A. JAIN, MD MFG:TEVA PLAN: PAIDMPD  
XXXI / KMN/KMN GROUP# CMD3896  
CLAIM REF# FKRWQHL

**Walgreens**  
12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

WED 1:30PM  
Refill

AZITHROMYCIN 250MG TABLETS  
00093-7146-56  
ALPHA

QTY 30  
10 DRAM



PINK  
FRONT: 93  
BACK: 7146

XXXI / KMN/KMN

Brunsting004465

AABrunsting.Financials003958

Ask if an antibiotic  
is causing her  
diaper rash.

*Without leaving her side.*



## New Pharmacy Chat at Walgreens.com

helps you find immediate answers anytime, anywhere.

Chat live with an expert from  
our pharmacy team about:

- Your personal prescription questions
- Over-the-counter product and prescription interactions
- Treatments for common ailments

Go to  
[Walgreens.com/pharmacychat](http://Walgreens.com/pharmacychat)  
today.  
It's confidential and secure!

*Walgreens*

Arm yourself®  
for the ones you love



No  
appointment  
necessary.

## Flu shots\* every day.

Register for your flu shot today  
at [Walgreens.com/flu](http://Walgreens.com/flu).

We bill many insurance plans directly,  
including Medicare.

\*Prices may vary for certain forms of the flu vaccine. Vaccines subject to availability. State, age and health condition-related restrictions may apply. See pharmacy for details.

*Walgreens*

8  
BR



NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

- Store TT 3 of 6
- Your Insurance Saved You: \$190.79

WED 1:30PM  
\$5.00  
EXPRESS PAY

03/02/11  
Refill



## Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION  
ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- Auto Refills  
We'll automatically refill your prescription before  
it's due to run out. Sign up in the pharmacy.

*Walgreens*  
The Pharmacy America Trusts • Since 1901™


Visit us online at [Walgreens.com](http://Walgreens.com)

Thank you for choosing Walgreens!

Brunsting004466

# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
12850 Memorial Drive  
Houston, TX 77024  
(713)722-7247

<b>PATIENT</b>	NELVA BRUNSTING	<b>DOCTOR</b>	A. JAIN, MD	<b>DRUG DESCRIPTION</b>
<b>BIRTH DATE</b>	10/08/26			
<b>MEDICATION</b>	SERTRALINE 50MG TABLETS	<b>PATIENT ALLERGIES</b>		
<b>QUANTITY</b>	30			
<b>DIRECTIONS</b>	TAKE 1 TABLET BY MOUTH EVERY DAY			
				BLUE
				FRONT: G 4900
				BACK: 50MG

**INGREDIENT NAME:** SERTRALINE (SER-tra-leen)

**COMMON USES:** This medicine is a selective serotonin reuptake inhibitor (SSRI) used to treat depression, panic disorder, obsessive-compulsive disorder (OCD), post-traumatic stress disorder (PTSD), social anxiety disorder (social phobia), and a severe form of premenstrual syndrome called premenstrual dysphoric disorder (PMDD). It may also be used for other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** WARNING: Antidepressants may increase the risk of suicidal thoughts or actions in children, teenagers, and young adults. However, depression and certain other mental problems may also increase the risk of suicide. Talk with the patient's doctor to be sure that the benefits of using this medicine outweigh the risks. Family and caregivers must closely watch patients who take this medicine. It is important to keep in close contact with the patient's doctor. Tell the doctor right away if the patient has symptoms like worsened depression, suicidal thoughts, or changes in behavior. Discuss any questions with the patient's doctor. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking a fenfluramine derivative (eg, dexfenfluramine); nefazodone; pirozamide; sibutramine; or thioridazine. **DO NOT TAKE THIS MEDICINE** if you are taking or have taken a monoamine oxidase inhibitor (MAOI) (eg, phenelzine), selegiline, or St. John's wort within the last 14 days. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking anxiolytics (eg, phenazepam); linezolid; metoclopramide; serotonergic 5-HT1 receptor agonists (eg, sumatriptan); trazodone; anticoagulants (eg, warfarin); aspirin; nonsteroidal anti-inflammatory drugs (NSAIDs) (eg, ibuprofen); diuretics (eg, furosemide, hydrochlorothiazide); tramadol; phenothiazines (eg, chlorpromazine); carbamazepine; cyproheptadine; atipirazole; clozapine; digoxin; flecainide; lithium; phenytoin; propafenone; risperidone; tricyclic antidepressants (eg, amitriptyline); or valproate (eg, valproic acid). **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including if you have a history of seizures; heart problems; liver problems; stomach or bowel bleeding; metabolism problems; allergies; pregnancy; or breast-feeding. Tell your doctor if you or a family member has a history of bipolar disorder (manic-depression), other mental or mood problems, suicidal thoughts or attempts, or alcohol or substance abuse. Tell your doctor if you are dehydrated, have low blood sodium levels, drink alcohol, or if you will be having electroconvulsive therapy (ECT). Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. This medicine has a Medication Guide approved by the U.S. Food and Drug Administration. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. **TAKE THIS MEDICINE** by mouth with or without food. **STORE THIS MEDICINE** at 75 degrees F (25 degrees C) away from heat, moisture, and light. Brief storage at temperatures between 59 and 86 degrees F (15 and 30 degrees C) is permitted. Do not store in the bathroom. **CONTINUE TO TAKE THIS MEDICINE** even if you feel well. Do not miss any doses. Taking this medicine at the same time each day will help you remember to take it. **DO NOT SUDDENLY STOP TAKING THIS MEDICINE** without checking with your doctor. Side effects may occur. They may include mental or mood changes, numbness or tingling of the skin, dizziness, confusion, headache, trouble sleeping, or unusual tiredness. You will be closely monitored when you start this medicine and whenever a change in dose is made. If you miss a dose of this medicine, take as soon as possible, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **DO NOT** take 2 doses at once.

**CAUTIONS:** **DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine. **THIS MEDICINE MAY CAUSE DROWSINESS OR DIZZINESS.** These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. **DO NOT DRIVE OR PERFORM OTHER POSSIBLY UNSAFE TASKS** until you know how you react to it. **DO NOT DRINK ALCOHOL** while you are taking this medicine. Check with your doctor before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers) while you are taking this medicine; they may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. Several weeks may pass before your symptoms improve. **DO NOT TAKE MORE THAN THE COMMANDED DOSE,** change your dose, or use this medicine for longer than prescribed without checking with your doctor. **IF YOUR DOCTOR**

**TELLS YOU TO STOP TAKING THIS MEDICINE,** you will need to wait for several weeks before beginning to take certain other medicines (eg, MAOIs, nefazodone). Ask your doctor when you should start to take your new medicines after you have stopped taking this medicine. **SEROTONIN SYNDROME** and **NEUROLEPTIC MALIGNANT SYNDROME (NMS)** are possibly fatal syndromes that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, triptans, MAOIs, antipsychotics). Symptoms of these syndromes may include blood pressure changes; agitation; confusion; hallucinations; other mental or mood changes; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; rigid muscles; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescribe or over-the-counter, check with your doctor or pharmacist. Caution is advised when using this medicine in the **ELDERLY;** they may be more sensitive to its effects, especially low blood sodium levels. Caution is advised when using this medicine in **CHILDREN;** they may be more sensitive to its effects, especially increased risk of suicidal thoughts or actions. **THIS MEDICINE MAY CAUSE WEIGHT CHANGES.** **CHILDREN AND TEENAGERS** may need regular weight and growth checks while they take this medicine. **FOR MEN,** THIS MEDICINE MAY RARELY CAUSE a prolonged, painful erection. This could happen even when you are not having sex. If this is not treated right away, it could lead to permanent sexual problems such as impotence. Contact your doctor right away if this happens. **FOR WOMEN,** THIS MEDICINE MAY CAUSE HARM TO THE FETUS if it is used during the last 3 months of pregnancy. **IF YOU BECOME PREGNANT,** contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. **IT IS NOT KNOWN IF THIS MEDICINE IS FOUND IN breast milk.** **IF YOU ARE OR WILL BE BREAST-FEEDING** while you use this medicine, check with your doctor. Discuss any possible risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include anxiety; constipation; decreased sexual desire or ability; diarrhea; dizziness; drowsiness; dry mouth; increased sweating; loss of appetite; nausea; nervousness; stomach upset; tiredness; trouble sleeping; vomiting; or weight loss. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience bizarre behavior; black or bloody stools; chest pain; decreased bladder control; exaggerated reflexes; fast or irregular heartbeat; fever; hallucinations; loss of coordination; new or worsening agitation, panic attacks, aggressiveness, impulsiveness, irritability, hostility, exaggerated feeling of well-being, restlessness, or inability to sit still; persistent or severe ringing in the ears; persistent, painful erection; red, swollen, blistered, or peeling skin; seizures; severe or persistent anxiety or trouble sleeping; stomach pain; suicidal thoughts or attempts; tremor; unusual bruising or bleeding; unusual or severe mental or mood changes; vision changes; or worsening of depression. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include coma; fainting; fast, slow, or irregular heartbeat; hair loss; hallucinations; seizures; severe or persistent dizziness, drowsiness, diarrhea, nausea, or vomiting; or tremor.

**ADDITIONAL INFORMATION:** **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** obtain refills before your supply runs out.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain. WIC# 957918

**NELVA BRUNSTING**  
13630 Pinerock, Houston, TX 770797517  
(713)484-4321  
**RX # 1515376-03328**      DATE: 03/02/11

**SERTRALINE 50MG TABLETS**  
QTY: 30      1 REFILL BEFORE 02/02/12  
Refill      NDC: 59762-4900-05  
Retail Price: \$29.99      Your Insurance Saved You: \$24.99

\$ 5.00

A. JAIN, MD  
MFG: CMD3896  
XXX/KMN/KMN / KMN  
PLAN: PAIDMPD  
GROUP# CMD3896  
CLAIM REF# OXHXMT3

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Customer Receipt

**NELVA BRUNSTING**  
13630 Pinerock, Houston, TX 770797517  
(713)484-4321  
**RX # 1515376-03328**      DATE: 03/02/11

**SERTRALINE 50MG TABLETS**  
QTY: 30      1 REFILL BEFORE 02/02/12  
Refill      NDC: 59762-4900-05  
Retail Price: \$29.99      Your Insurance Saved You: \$24.99

\$ 5.00

A. JAIN, MD  
MFG: GREENSTONE  
XXX/KMN/KMN / KMN  
PLAN: PAIDMPD  
GROUP# CMD3896  
CLAIM REF# OXHXMT3

**Walgreens**

12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

WED 1:30PM  
Refill

SERTRALINE 50MG TABLETS  
59762-4900-05  
CELL 29

QTY 30  
20 DRAM



BLUE  
FRONT: G 4900  
BACK: 50MG

XXX/KMN/KMN / KMN

Med Guide

Brunsting004463

AABrunsting.Financials003960

Ask if grapefruit juice affects your medication.

From the breakfast table.



# New Pharmacy Chat at Walgreens.com

helps you find immediate answers anytime, anywhere.

Chat live with an expert from our pharmacy team about:

- Your personal prescription questions
- Over-the-counter product and prescription interactions
- Treatments for common ailments




Go to [Walgreens.com/pharmacychat](http://Walgreens.com/pharmacychat) today.

It's confidential and secure!

*Walgreens*

## Arm yourself® for the ones you love



No appointment necessary.

# Flu shots\* every day.

Register for your flu shot today at [Walgreens.com/flu](http://Walgreens.com/flu).

We bill many insurance plans directly, including Medicare.

\*Prices may vary for certain forms of the flu vaccine. Vaccines subject to availability. State, age and health condition-related restrictions may apply. See pharmacy for details.

*Walgreens*

8 BR



NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

- Store TT 5 of 6
- Your Insurance Saved You: \$24.99

WED 1:30PM  
\$5.00  
EXPRESS PAY

03/02/11  
Refill  
MED GUIDE



## Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- Auto Refills  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

*Walgreens*  
The Pharmacy America Trusts • Since 1901™

Visit us online at [Walgreens.com](http://Walgreens.com)

Thank you for choosing Walgreens!

Brunsting004464

# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
12850 Memorial Drive  
Houston, TX 77024  
(713)722-7247

<b>PATIENT</b>	NELVA BRUNSTING	<b>DOCTOR</b>	A. JAIN, MD	<b>DRUG DESCRIPTION</b>	LIQUID
<b>BIRTH DATE</b>	10/08/26				
<b>MEDICATION</b>	BROVANA 15MCG/2ML INH SOL 30X2ML	<b>PATIENT ALLERGIES</b>			
<b>QUANTITY</b>	120				
<b>DIRECTIONS</b>	INHALE 1 VIAL VIA NEBULIZER TWICE DAILY				

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** ARFORMOTEROL (ar-for-MOE-tar-cle)

**COMMON USES:** This medicine is a long-acting beta-agonist bronchodilator used for long-term treatment of chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE. WARNING: LONG-ACTING BETA-AGONISTS SUCH AS THIS MEDICINE HAVE BEEN RARELY ASSOCIATED WITH AN INCREASED RISK OF ASTHMA-RELATED DEATH.** Long-acting beta-agonists should not be used in asthma patients without another long-term asthma-control medicine (eg, inhaled corticosteroid). This medicine has not been approved to treat asthma. Safety and effectiveness of this medicine in patients with asthma have not been confirmed. **SOME MEDICINES MAY INTERACT** with this medicine. **DO NOT TAKE THIS MEDICINE** if you are using another medicine that has a long-acting beta-agonist (eg, tiotropium) or if you are taking any of the following: corticosteroids (eg, prednisone), diuretics (eg, furosemide, hydrochlorothiazide), xanthines (eg, theophylline), cardiac medications (eg, digoxin), tricyclic antidepressants (eg, amitriptyline, nortriptyline), monoamine oxidase inhibitors (MAOIs) (eg, phenelzine), tricyclic antidepressants (eg, amitriptyline, nortriptyline), and others. **TALK WITH YOUR DOCTOR** for a complete list of all interactions that may occur. Ask your health care provider if this medicine may interact with other medicines you take. **DO NOT START OR STOP any medicine without your doctor's approval.** Inform your doctor of any other medical conditions, including a history of other breathing problems (eg, asthma), diabetes, heart problems (eg, fast or irregular heartbeat, heart blood vessel problems), liver problems, high blood pressure, low blood potassium levels, seizures, an overactive thyroid, or if you have high blood or urine ketone levels. **Inform your doctor if you have any of the following:** heart failure, allergies, pregnancy, or breast-feeding. Tell your doctor if you have recently been to an emergency room for breathing problems; have a history of frequent hospitalizations for breathing problems; have ever had life-threatening breathing problems; or have had an unusual reaction to a sympathomimetic medicine (eg, albuterol, pseudoephedrine), such as fast or irregular heartbeat, overexcitement, or severe trouble sleeping. Tell your doctor if you have taken a MAOI (eg, phenelzine) or a tricyclic antidepressant (eg, amitriptyline) within the last 14 days. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you are having severe breathing problems (eg, sudden, severe onset or worsening of COPD symptoms such as chest tightness, cough, shortness of breath, wheezing); you have asthma and you are not currently using a long-term asthma-control medicine (eg, inhaled corticosteroids); or if you have asthma that is severe and well-controlled with the use of a long-term asthma-control medicine. **THIS MEDICINE SHOULD NOT BE USED IN CHILDREN;** safety and effectiveness in children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine comes with a **MEDICATION GUIDE** approved by the US Food and Drug Administration. Read it carefully each time you refill this medicine. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. **THIS MEDICINE SHOULD ONLY BE USED WITH A NEBULIZER.** Do NOT inject or swallow it. A health care provider will teach you how to use the nebulizer. Be sure you know what type of nebulizer to use with this medicine and how to use it. Follow the procedures you are taught when you use a dose. Contact your health care provider if you have any questions. **STORE THIS MEDICINE** in the original foil pouch and get it out of the pouch from the foil pouch or the vial until right before use. **IF THE MEDICINE CONTAINS PARTICLES,** is cloudy or discolored, or if the vial is cracked or damaged in any way, do not use it. **DO NOT MIX THIS MEDICINE WITH OTHER MEDICINES** in your nebulizer machine. **TO USE THIS MEDICINE,** twist open the top of the vial and pour the entire contents into the nebulizer reservoir. **CONNECT THE NEBULIZER RESERVOIR** to the mouthpiece or face mask. **CONNECT THE NEBULIZER** to the compressor. Sit in a comfortable, upright position. **PLACE THE MOUTHPIECE** in your mouth or put on the face mask and turn on the nebulizer. **BREATHE AS CALMLY,** deeply, and evenly as possible until no more mist is formed in the nebulizer chamber (about 5 to 10 minutes). **USE THE NEBULIZER** according to the instructions. Failure to properly clean the nebulizer could lead to bacteria entering the medicine. This may lead to an infection. To avoid bacteria entering the medicine, use the entire contents right after opening the vial for the first time. **STORE THIS MEDICINE** in the REFRIGERATOR, between 36 and 46 degrees Fahrenheit (2 and 10 degrees Celsius). Do not freeze. **UNOPENED POUCHES** MAY BE STORED at room temperature between 68 and 77 degrees Fahrenheit (20 and 25 degrees Celsius) for no more than 6 weeks from the date of manufacture and light. If this medicine is stored at room temperature, throw it away after 6 weeks. **DO NOT USE** this medicine if it is past the expiration date on the container. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **CONTINUE TO USE THIS MEDICINE** even if you feel well. Do not miss any doses. If you miss a dose of the medicine, skip the missed dose and go back to your regular dosing schedule. Do not use 2 doses at once.

**CAUTIONS: DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine or to formoterol. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescription or over-the-counter, check with your doctor or pharmacist. **LAB TESTS,** including lung function and blood potassium levels, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to attend all doctor and lab appointments. **THIS MEDICINE MAY CAUSE DIZZINESS.** This effect may be worse if you take it with alcohol or other medicines. Use this medicine with caution. Do not drive or perform other possibly unsafe tasks until you know how you react to it. **THIS MEDICINE DOES NOT STOP SUDDEN SYMPTOMS OF COPD ONCE THEY HAVE ALREADY STARTED.** Be sure to carry a short-acting bronchodilator inhaler (eg, albuterol) with you at all times to treat any breathing problems that may occur between doses of this medicine (eg, severe or sudden onset of wheezing or shortness of breath). If you have any questions about which medicines stop sudden symptoms, check with your doctor or pharmacist. **IF YOU HAVE BEEN PREVIOUSLY USING A SHORT-ACTING BRONCHODILATOR INHALER,** talk with your doctor about how to use both this medicine. Short-acting bronchodilators are normally used with this medicine to treat breathing problems that may occur between doses. **THE RISK OF SERIOUS HEART PROBLEMS** (eg, irregular heartbeat) may be greater if you use this medicine in high doses. Do NOT use more than recommended dose or use more often than prescribed. **TELL YOUR DOCTOR AT ONCE** if you notice that your short-acting bronchodilator inhaler does not work as well as you need to use it often, or if your breathing problems get worse. **CONTACT YOUR DOCTOR OR SEEK MEDICAL CARE RIGHT AWAY** if you have breathing problems that worsen quickly, or if you use your short-acting bronchodilator and do not get relief. **TALK WITH YOUR DOCTOR OR PHARMACIST** about all of your breathing medicines and how to use them. Do not start, stop, or change the dose of any

breathing medicine unless your doctor tells you to. **THE MEDICINE MAY CAUSE SERIOUS BREATHING PROBLEMS RIGHT AFTER YOU USE A DOSE.** If this happens, use your short-acting bronchodilator. Contact your doctor or seek other medical care at once. **THIS MEDICINE MAY RAISE YOUR BLOOD SUGAR.** High blood sugar may make you feel confused, drowsy, or thirsty. It can also make you flush, breathe faster, or have a fruit-like breath odor. If these symptoms occur, tell your doctor right away. **PREGNANCY AND BREAST-FEEDING:** If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. It is not known if this medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include: back pain; diarrhea; dry mouth; headache; nausea; nervousness; stuffy nose; tiredness; tremor; trouble sleeping; vomiting. If they continue or are bothersome, check with your doctor. **SEEK MEDICAL ATTENTION RIGHT AWAY** if any of these SEVERE side effects occur: chest pain; fast or irregular heartbeat; fever, chills, or persistent sore throat; leg swelling; new or worsening breathing problems (eg, increased chest tightness, coughing, shortness of breath, wheezing); severe or persistent headache, dizziness, tremor, or nervousness; severe or persistent muscle pain or cramps; symptoms of high blood sugar (eg, increased thirst, urination, or hunger; unusual weakness or drowsiness; confusion); trouble speaking. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, tongue, or throat; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE: IF OVERDOSE IS SUSPECTED,** contact your local poison control center or emergency room immediately. Symptoms may include chest pain; fast or irregular heartbeat; severe or persistent dizziness, dry mouth, fatigue, headache, muscle pain or cramps, nausea, nervousness, trouble sleeping, or tremors; severe or persistent symptoms of high blood sugar (eg, increased thirst, urination, or hunger; drowsiness; flushing of the skin; confusion; fruity-taste breath odor).

**ADDITIONAL INFORMATION:** If your symptoms do not improve or if they become worse, check with your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD** OR IF YOU obtain refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN. STORE IN SAFETY CONTAINER OR SECURE AREA.**

**NELVA BRUNSTING**  
13630 Pinerock, Houston, TX 770797517  
(713)464-4391

**RX # 1540088-03328** DATE: 04/21/11

**BROVANA 15MCG/2ML INH SOL 30X2ML**  
QTY: 120 2 REFILLS BEFORE 04/21/12  
Copy NDC: 63402-0911-30 \$ 527.99

A. JAIN, MD  
MFG: SEPRACOR  
XXX/RJW/KHT /PBP

**Walgreens** 12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247 Customer Receipt

**NELVA BRUNSTING**  
13630 Pinerock, Houston, TX 770797517  
(713)464-4391

**RX # 1540088-03328** DATE: 04/21/11

**BROVANA 15MCG/2ML INH SOL 30X2ML**  
QTY: 120 2 REFILLS BEFORE 04/21/12  
Copy NDC: 63402-0911-30 \$ 527.99

A. JAIN, MD  
MFG: SEPRACOR  
XXX/RJW/KHT /PBP

**Walgreens** 12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247 Duplicate Receipt

Pharmacy use only

THU 5:16PM	BROVANA 15MCG/2ML INH SOL 30X2ML	QTY 120	LIQUID
Copy	REFRIG		
		XXX/RJW/KHT /PBP	Med Guide

Brnunsting004457

AABrunsting.Financials003962

Do not flush unused medications or pour down a sink or drain. WIC# 957918

# Save up to 30% on your prescriptions

Enroll today for only \$19.95!

The AARP® Prescription Discount Program from Walgreens saves you up to 30 percent on all your FDA-approved prescriptions. This includes generic, brand name and specialty medications.



## How Does It Work?

Your AARP Prescription Discount Program card can be used for drugs not covered by your insurance plan or if you have no prescription insurance. Simply present your AARP Prescription Discount Program card to your Walgreens pharmacist — after any other insurance card you may also have — when you fill or refill a prescription.

Note: This program is not a prescription drug insurance plan or a Medicare Part D plan, and does not replace such coverage.

## How Do I Enroll?

You must be an AARP member to enroll. We can enroll you into both programs with one easy call. Just dial 1-877-4AARP19 (1-877-422-7719). You can also enroll online by visiting us at [www.aarp-pharmacy.com](http://www.aarp-pharmacy.com).

## What If I Have Questions?

Visit [aarp-pharmacy.com](http://aarp-pharmacy.com) or call the Walgreens Customer Care Center toll free, 24/7 at 1-877-4AARP19 (1-877-422-7719), or TTY 1-800-925-0178.

8  
BR



\*1540088 0801 3 0052799 5\*

NELVA BRUNSTING

13630 Pinerock  
Houston, TX 770797517  
(713)464-4381

- PAIDMP: Prior Authorization Required
- Prescription Savings Club could save you \$117.29!
- Ask if you qualify.

THU 5:16PM  
\$527.99  
EXPRESS PAY

04/21/11  
Copy

REFRIGERATE  
MED GUIDE



## Personal Prescription Information

### LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.

Take advantage of these convenient services:

- **Touch Tone Refills**  
Save time by using our automated system for a refill. Just dial the number on your prescription label.
- **Auto Refills**  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

**Walgreens**  
The Pharmacy America Trusts • Since 1901™

Visit us online at [Walgreens.com](http://Walgreens.com)

**Thank you for choosing Walgreens!**

OFF: 3875 677 116 0935993

INFC: 0916 63402091130

57616

AA Brunsting Financials 003963

Brunsting004458




# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
12850 Memorial Drive  
Houston, TX 77024  
(713)722-7247

<p><b>PATIENT</b> NELVA BRUNSTING  <b>BIRTH DATE</b> 10/08/26  <b>MEDICATION</b> ETHAMBUTOL 400MG TABLETS  <b>QUANTITY</b> 90  <b>DIRECTIONS</b> TAKE 3 TABLETS BY MOUTH EVERY DAY</p>	<p><b>DOCTOR</b> A. JAIN, MD  <b>PATIENT ALLERGIES</b></p>
--	--

**DRUG DESCRIPTION**



WHITE  
FRONT: L U  
BACK: C32

---

**INGREDIENT NAME:** ETHAMBUTOL (e-THAM-byoo-tole)

**COMMON USES:** This medicine is an antibacterial used to treat tuberculosis (TB).

**BEFORE USING THIS MEDICINE:** INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. Inform your doctor of any other medical conditions, allergies, pregnancy, or breast-feeding.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. STORE THIS MEDICINE at room temperature in a tightly-closed container, away from heat and light. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** DO NOT STOP USING THIS MEDICINE without first checking with your doctor. THIS MEDICINE MAY CAUSE dizziness. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. HAVE REGULAR EYE EXAMINATIONS while you are taking this medicine even if you do not notice changes in your vision. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS, that may go away during

**ADDITIONAL INFORMATION:** treatment, include nausea, headache, or dizziness. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience changes in vision, vomiting, skin rash, itching, fever, confusion, or numbness or tingling of extremities. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately.

**ADDITIONAL INFORMATION:** If your symptoms do not improve within 2 to 3 weeks, or if they become worse, check with your doctor. DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

**NELVA BRUNSTING**  
13630 Pinecroft, Houston, TX 770797617  
(713)464-4391  
**RX # 1534700-03328**      DATE: 04/29/11

---

**ETHAMBUTOL 400MG TABLETS**  
QTY: 90      1 REFILL BEFORE 04/05/12  
Refill      NDC: 68180-0281-01  
Retail Price: \$153.59      Your Insurance Saved You: \$148.59      \$ 5.00

A. JAIN, MD      PLAN: PAIDMPP  
MFG: LUPIN      GROUP# CMD3896  
XXX/PBP/PBP/ /PBP      CLAIM REF# 7C1HH79

**Walgreens**      12850 MEMORIAL DRIVE HOUSTON, TX 77024      PH: (713)722-7247      Customer Receipt

**NELVA BRUNSTING**  
13630 Pinecroft, Houston, TX 770797617  
(713)464-4391  
**RX # 1534700-03328**      DATE: 04/29/11


---

**ETHAMBUTOL 400MG TABLETS**  
QTY: 90      1 REFILL BEFORE 04/05/12  
Refill      NDC: 68180-0281-01  
Retail Price: \$153.59      Your Insurance Saved You: \$148.59      \$ 5.00

A. JAIN, MD      PLAN: PAIDMPP  
MFG: LUPIN      GROUP# CMD3896  
XXX/PBP/PBP/ /PBP      CLAIM REF# 7C1HH79

**Walgreens**      12850 MEMORIAL DRIVE HOUSTON, TX 77024      PH: (713)722-7247      Duplicate Receipt

Pharmacy use only

SUN 11:00AM Refill	ETHAMBUTOL 400MG TABLETS 68180-0281-01 ALPHA	QTY 90		WHITE FRONT: L U BACK: C32
--------------------	--	--------	---	----------------------------------

XXX/PBP/PBP/ /PBP

Brunsting004461



## Take your antibiotics the right way.

- Precisely follow usage directions.
- Do not skip doses.
- Do not share them with others.
- Finish the prescription even if you feel better.
- Do not save them for future use.

### Why is this checklist so important?

Using an antibiotic the wrong way can make infections stronger and harder to treat. You can prevent this problem by getting smart about antibiotics.

For more information talk to your Walgreens pharmacist. Or call 1-800-CDC-INFO or visit [www.cdc.gov/getsmart](http://www.cdc.gov/getsmart).

5785

## Do you have a higher risk of getting pneumonia?

### Are you:

- Age 19 through 64 and smoke or have asthma?
- Age 64 or younger and have diabetes, heart disease, lung disease, leukemia, lymphoma, Hodgkin's disease, kidney problems, HIV or other condition that lowers the body's resistance to infection?
- Age 64 or younger and are taking a drug or treatment that lowers the body's resistance to infection, such as long-term steroids, certain cancer drugs or radiation therapy?
- Age 65 or older?

If you answered, "yes" to any of these questions, the Centers for Disease Control & Prevention (CDC) recommends that you receive a pneumonia vaccination.

**Talk to your Walgreens pharmacist to get vaccinated today!**

No out-of-pocket cost for Medicare Part B beneficiaries\*

There's a way to stay well.

\*Medicare Part B generally covers the pneumonia vaccine once per beneficiary. Vaccine subject to availability. State, age and health condition-related restrictions may apply. See pharmacy for details.

OFF: 2875 677 119 0064305

INFO: 0917 68130028101

5785

8  
BR




PAIDMPD \*1534700 0302 3 0000500 1\*

NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

- Store TT 1 of 5
- Your Insurance Saved You: \$148.59

SUN 11:00AM  
\$5.00  
EXPRESS PAY

04/29/11  
Refill



**Personal Prescription Information**

### LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- Auto Refills  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

The Pharmacy America Trusts • Since 1801™


Visit us online at [Walgreens.com](http://Walgreens.com)

**Thank you for choosing Walgreens!**

Brunsting004462

# YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
12850 Memorial Drive  
Houston, TX 77024  
(713)722-7247

<b>PATIENT</b>	NELVA BRUNSTING	<b>DOCTOR</b>	A. JAIN, MD	<b>DRUG DESCRIPTION</b>
<b>BIRTH DATE</b>	10/08/26			 PINK FRONT: m 32
<b>MEDICATION</b>	METOPROLOL TARTRATE 50MG TABLETS	<b>PATIENT ALLERGIES</b>		
<b>QUANTITY</b>	60			
<b>DIRECTIONS</b>	TAKE 1 TABLET BY MOUTH EVERY 12 HOURS			

**INGREDIENT NAME:** METOPROLOL (me-TOE-proe-lole)

**COMMON USES:** This medicine is a beta-adrenergic blocking agent (beta-blocker) used to treat high blood pressure. It may be used alone or with other medicines. It is also used for the long-term treatment of chest pain (angina) and to reduce the risk of death due to heart problems in certain patients who have had a heart attack. It may also be used for other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** **WARNING:** Do not suddenly stop taking this medicine; sharp chest pain, irregular heartbeat, and sometimes heart attack may occur. The risk may be greater if you have certain types of heart disease. Your doctor should slowly lower your dose over several weeks if you need to stop taking it, even if you only take it for high blood pressure. Heart disease is common and you may not know you have it. Do not suddenly stop taking this medicine before you receive any medical or dental care, emergency care, or surgery. Tell your doctor if you have a history of diabetes or take medicine to lower your blood sugar (eg, glyburide, insulin). This medicine may hide signs of low blood sugar such as fast heartbeat. Tell your doctor right away if you notice other signs of low blood sugar such as anxiety, chills, dizziness, drowsiness, fainting, headache, tremor, unusual sweating, vision changes, or weakness. This medicine should not usually be used by patients who have a history of certain lung or breathing problems (eg, asthma) or a certain type of adrenal gland tumor (pheochromocytoma). It may worsen these conditions. Tell your doctor if you have a history of these conditions; your doctor may need to adjust your dose or prescribe additional medicine to reduce the risk of side effects. Tell your doctor if you have a history of overactive thyroid. This medicine may hide symptoms of overactive thyroid (eg, fast heartbeat). Do not suddenly stop taking this medicine; suddenly stopping this medicine could worsen your condition. Your doctor should slowly lower your dose over several weeks if you need to stop taking it. Check with your doctor for more information. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking mifepristol. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking amiodarone, bupropion, certain HIV protease inhibitors (eg, ritonavir), certain selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), cimetidine, digoxin, difenhydramine, disopyramide, flecainide, hormonal contraceptives (eg, birth control pills), hydralazine, hydroxychloroquine, ketanserin, mofloquine, phenothiazines (eg, thioridazine), propafenone, quazolidines (eg, alfuzosin), quinine, terbinafine, propylthiouracil, verapamil, barbiturates (eg, phenobarbital), indomethacin, phenylpropranolamine, bupivacaine, lidocaine, or clonidine. **DO NOT START OR STOP** any medicine without your doctor or pharmacist approval. Inform your doctor of any other medical conditions including lung or breathing problems (eg, asthma, bronchitis, chronic obstructive pulmonary disease [COPD]), diabetes, low blood pressure, thyroid problems, or adrenal gland tumor (pheochromocytoma), allergies, pregnancy or breastfeeding. Tell your doctor if you have a history of heart attack, slow or irregular heartbeat, heart failure, or other heart problems; chest pain or angina; blood vessel or circulation problems; liver disease; or if you will be having surgery or receiving anesthesia. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have a very slow heartbeat (eg, bradycardia), certain types of irregular heartbeat (eg, atrioventricular [AV] block, sick sinus syndrome), moderate to severe heart failure, very low systolic blood pressure (less than 100 mm Hg), or severe blood circulation problems. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. **TAKE THIS MEDICINE BY MOUTH WITH FOOD** or immediately following a meal at the same time each day. **STORE THIS MEDICINE** at room temperature, between 59 and 86 degrees F (15 and 30 degrees C). Store away from heat, moisture, and light. Do not store in the bathroom. **IF YOU MISS A DOSE OF THIS MEDICINE,** take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** **DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine or to another beta-blocker (eg, propranolol). **IF YOU HAVE A HISTORY OF ANY SEVERE ALLERGIC REACTION,** talk with your doctor. You may be at risk for an even more severe allergic reaction if you ever do contact with the substance that caused your allergy. Some medicines used to treat

severe allergies may also not work as well while you are using this medicine. **LAB TESTS,** including liver and kidney function, blood pressure, and complete blood cell counts, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to keep all doctor and lab appointments. **THIS MEDICINE MAY CAUSE DROWSINESS, DIZZINESS, OR LIGHTEADEDNESS.** These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. Do not drive or perform other possibly unsafe tasks until you know how you react to it. This medicine may cause dizziness, lightheadedness, or fainting; alcohol, hot weather, exercise, or fever may increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of any of these effects. **PATIENTS WHO TAKE MEDICINE FOR HIGH BLOOD PRESSURE** often feel tired or run down for a few weeks after starting treatment. Be sure to take your medicine even if you may not feel "normal." Tell your doctor if you develop any new symptoms. **DO NOT SUDDENLY STOP TAKING THIS MEDICINE** without first consulting your doctor. If your doctor decides you should no longer use this medicine, you will need to stop this medicine gradually according to your doctor's instructions. **IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK YOUR BLOOD PRESSURE** and heart rate regularly, be sure to do so. **TELL YOUR DOCTOR OR DENTIST** that you take this medicine before you receive any medical or dental care, emergency care, or surgery. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescription or over-the-counter, check with your doctor or pharmacist. Do not take any medicines used to treat colds or congestion without first talking with your doctor or pharmacist. **FOR WOMEN:** If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. This medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby. **DIABETES PATIENTS:** this medicine may hide signs of low blood sugar, such as rapid heartbeat. Be sure to watch for other signs or low blood sugar. Low blood sugar may make you anxious, sweaty, weak, dizzy, drowsy, or faint. It may also make your vision change; give you a headache, chills, or tremors; or make you more hungry. Check blood sugar levels closely. Ask your doctor before you change the dose of your diabetes medicine.

**POSSIBLE SIDE EFFECTS:** **SIDE EFFECTS** that may occur while taking this medicine include constipation, diarrhea, dizziness, dry mouth/eyes, gas, headache, heartburn, lightheadedness, mild drowsiness, muscle aches, nausea, stomach pain, trouble sleeping, unusual tiredness or weakness, or vomiting. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience blue or unusually cold hands or feet; chest pain; fainting; hallucinations; mood or mental changes (eg, confusion, depression); pounding in the chest; severe dizziness or lightheadedness; shortness of breath; slow or irregular heartbeat; swelling of the arms, hands, and feet; vision changes; wheezing; yellowing of the skin or eyes. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include chest pain, seizures, very slow heart beat, severe dizziness, fainting, and difficult or slowed breathing.

**ADDITIONAL INFORMATION:** **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** obtain refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain. WIC# 957918

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**NELVA BRUNSTING**  
13830 Pinerock, Houston, TX 770797517  
(713)464-4391  
**RX # 1534699-03328**      DATE: 04/29/11

**METOPROLOL TARTRATE 50MG TABLETS**  
QTY: 60    1 REFILL BEFORE 04/05/12  
Refill    NDC:00378-0032-10  
Retail Price: \$16.66    Your Insurance Saved You: \$12.28    \$ 4.38

A. JAIN, MD                      PLAN: PAIDMFD  
MFG:MYLAN                  GROUP# CMD3896  
XXX/PBP/PBP /PBP        CLAIM REF# SLOMDCX

**Walgreens**      12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247      Customer Receipt

Pharmacy use only

SUN 11:00AM      METOPROLOL TARTRATE 50MG TABLETS      QTY 60  
Refill                      00378-0032-10                      20 DRAM  
CELL 138                      FRONT: m 32

**NELVA BRUNSTING**  
13830 Pinerock, Houston, TX 770797517  
(713)464-4391  
**RX # 1534699-03328**      DATE: 04/29/11

**METOPROLOL TARTRATE 50MG TABLETS**  
QTY: 60    1 REFILL BEFORE 04/05/12  
Refill    NDC:00378-0032-10  
Retail Price: \$16.66    Your Insurance Saved You: \$12.28    \$ 4.38

A. JAIN, MD                      PLAN: PAIDMFD  
MFG:MYLAN                  GROUP# CMD3896  
XXX/PBP/PBP /PBP        CLAIM REF# SLOMDCX

**Walgreens**      12850 MEMORIAL DRIVE HOUSTON, TX 77024  
PH: (713)722-7247      Duplicate Receipt

 PINK  
FRONT: m 32

XXX/PBP/PBP /PBP

Brunsting004459

AABrunsting.Financials003966



# Cold or flu? Antibiotics aren't for you.

Antibiotics kill bacteria,  
not viruses.

- Antibiotics can cure most bacterial infections, such as sore throats caused by strep and bacterial sinus infections.
- Using antibiotics for viral illness, like the common cold, will not help you feel better or prevent spreading it.

Please follow your healthcare provider's advice. And to learn more about antibiotics, visit [www.cdc.gov/getsmart](http://www.cdc.gov/getsmart) today.

5896



# When you're sick, antibiotics aren't always the answer.

To avoid antibiotic-resistant infections and adverse drug events, avoid seeking an antibiotic prescription for colds, coughs and sniffles.

- Taking antibiotics when they are not needed causes some bacteria to become resistant to the antibiotic.
- Resistant bacteria are stronger and make future bacterial illnesses harder to treat.

To learn more, talk with your  
Walgreens pharmacist today.

OPT-2875 677 119 0963941

INRC-0917 00378003210

5896

8  
BR




PAIDMPD \*1534699 0302 3 0000438 5\*

NELVA BRUNSTING  
13630 Pinerock  
Houston, TX 770797517  
(713)464-4391

- Store TT 2 of 5
- Your Insurance Saved You: \$12.28

SUN 11:00AM  
\$4.38  
EXPRESS PAY

04/29/11  
Refill



**Personal  
Prescription  
Information**

**LOOK INSIDE FOR IMPORTANT INFORMATION  
ABOUT YOUR MEDICATION.**

Take advantage of this convenient service:

- Auto Refills  
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

**Walgreens**  
The Pharmacy America Trusts • Since 1901™

Visit us online at [Walgreens.com](http://Walgreens.com)

**Thank you for choosing Walgreens!**

Brunsting004460