

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
12850 Memorial Drive
Houston, TX 77024
(713)722-7247

PATIENT NELVA BRUNSTING **DOCTOR** A. JAIN, MD **DRUG DESCRIPTION**
BIRTH DATE 10/08/26
MEDICATION SPIRIVA CAPS 30'S & HANDIHALER
QUANTITY 30 **PATIENT ALLERGIES**
DIRECTIONS INHALE CONTENTS OF ONE CAPSULE ONCE DAILY USING HANDIHALER

G Losen

INGREDIENT NAME: TIOTROPIUM BROMIDE (TYE-oh-TROE-pee-um BROE-mide)

COMMON USES: This medicine is an anticholinergic agent used to treat the symptoms of chronic obstructive pulmonary disease, including chronic bronchitis and emphysema. It may also be used to treat other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking another anticholinergic (eg, ipratropium). Ask your doctor if you are unsure if any of your medicines are anticholinergics. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including if you have trouble urinating; an enlarged prostate; bladder blockage; glaucoma; kidney problems; allergy (to milk proteins); pregnancy; or breast-feeding. Tell your doctor if you are having an asthma attack or increased difficulty breathing. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. An additional patient information leaflet is available with this medicine. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have before using this medicine. **DO NOT SWALLOW THE CAPSULES. THE CAPSULES ARE USED WITH A SPECIAL INHALER.** **DO NOT** remove a capsule from the packaging until you are ready to use it. To remove a capsule from the packaging, carefully peel the foil back to expose the capsule, then tip the capsule out of the blister. **DO NOT** cut the foil or use sharp objects to remove the capsule from the blister. If a second capsule is exposed to the air when you are removing a capsule for use, it must be discarded. Do not save the capsule for later. Place the capsule in the special inhaler device immediately. The device will puncture the capsule so that the medicine inside may be inhaled into the lungs through the mouthpiece. Exhale slowly and deeply. **DO NOT** breathe into the mouthpiece of the inhaler. Position the inhaler mouthpiece between your lips and try to rest your tongue flat. Keep your head upright. Do not block the air vents on the inhaler. **TAKE A SLOW, DEEP BREATH.** You should hear or feel the capsule vibrate inside the inhaler. Hold your breath as long as it feels comfortable, then exhale slowly through pursed lips. Breathe out completely. To be sure all of the medicine has been inhaled from the capsule, inhale from the mouthpiece a second time following the same process. **ASK YOUR DOCTOR OR PHARMACIST** if you are unclear on how to use this device or inhale the medicine. **CLEAN THE INHALER DEVICE** once a month according to the instructions in the patient leaflet. Each inhaler device may be used for up to 1 year and then should be replaced. **STORE THIS MEDICINE** at room temperature at 77 degrees F (25 degrees C), away from heat, moisture, and light. Brief storage between 59 and 86 degrees F (15 and 30 degrees C) is permitted. Do not store in the bathroom. This medicine will work best if used regularly. Taking your dose at the same time each day will help you to remember. **IF YOU MISS A DOSE OF THIS MEDICINE,** use it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **DO NOT** use 2 doses at once. **DO NOT** use this medicine more often than 1 time every 24 hours.

CAUTIONS: **DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine or to atropine or related medicines (eg, ipratropium). **THIS MEDICINE WILL NOT STOP AN ASTHMA ATTACK** once one has started. **IF YOU ARE ALSO USING A RESCUE INHALER** (eg, albuterol), be sure to always carry the bronchodilator inhaler with you to use during asthma attacks. **DO NOT EXCEED THE RECOMMENDED DOSE** without checking with your doctor. **DO NOT STOP USING THIS MEDICINE** without first checking with your doctor. This medicine may cause dizziness or blurred vision. These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. **AVOID GETTING THIS MEDICINE IN YOUR EYES.** If you get the medicine in your eyes and eye pain, blurred vision, or other vision changes occur, contact your doctor immediately. **THIS MEDICINE MAY SOMETIMES CAUSE SEVERE BREATHING PROBLEMS** right after you use a dose. If this happens, use your short-acting bronchodilator inhaler. Contact your doctor or seek other medical care at once. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE,** either prescription or over-the-counter, check with your doctor or pharmacist. **USE THIS MEDICINE WITH CAUTION IN THE ELDERLY;** they may be more sensitive to its effects, especially constipation and urinary tract infections. **FOR WOMEN:** If you

PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. **IT IS NOT KNOWN IF THIS MEDICINE IS FOUND** in breast milk. **IF YOU ARE OR WILL BE BREAST-FEEDING** while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

POSSIBLE SIDE EFFECTS: **SIDE EFFECTS** that may occur while using this medicine include dry mouth, constipation, sinus inflammation, throat irritation, stomach pain, vomiting, blurred vision, or mild nosebleed. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience irritation, pain, or white patches in your mouth or on your tongue; mouth sores; severe or persistent nosebleeds; severe or persistent constipation; difficult or painful urination; eye pain or discomfort; vision changes (eg, halos, colored images); chest pain; fast or irregular heartbeat; difficulty swallowing; new or worsened breathing problems; or wheezing. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: **IF OVERDOSE IS SUSPECTED,** contact your local poison control center or emergency room immediately. Symptoms may include mental changes; severe constipation; stomach pain; or tremors.

ADDITIONAL INFORMATION: **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** obtain refills before your supply runs out.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

NELVA BRUNSTING

13630 Pinrock, Houston, TX 770797517
(713)464-4391

RX # 1496586-03328 **DATE:** 12/05/10

SPIRIVA CAPS 30'S & HANDIHALER

QTY: 30 **3 REFILLS BEFORE 12/05/11**
New **NDC:** 00597-0075-41
Retail Price: \$236.99 **Your Insurance Saved You:** \$215.99

\$ 21.00

A. JAIN, MD
MFG:BOEHRINGER
TPL/RJW/RJW/RJW/RJW

PLAN: PAIDMPD
GROUP# CMD3896
CLAIM REF# RDC9TAQ

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
PH: (713)722-7247

Customer Receipt

NELVA BRUNSTING

13630 Pinrock, Houston, TX 770797517
(713)464-4391

RX # 1496586-03328 **DATE:** 12/05/10

SPIRIVA CAPS 30'S & HANDIHALER

QTY: 30 **3 REFILLS BEFORE 12/05/11**
New **NDC:** 00597-0075-41
Retail Price: \$236.99 **Your Insurance Saved You:** \$215.99

\$ 21.00

A. JAIN, MD
MFG:BOEHRINGER
TPL/RJW/RJW/RJW/RJW

PLAN: PAIDMPD
GROUP# CMD3896
CLAIM REF# RDC9TAQ

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

SUN 1:29PM **SPIRIVA CAPS 30'S & HANDIHALER** **QTY 30**
New **ALPHA** **00597-0075-41**

TPL/RJW/RJW/RJW/RJW

Brunsting004489

AABrunsting.Financials003940

Get a **FREE** one-on-one
Medicare Part D
review session!

Your pharmacist will:

- Review your medications and look for ways to help save you money
- Talk about other things you can do to stay well
- Check for potential drug interactions



Make your appointment today!

Walgreens

There's a way to stay well.

Information provided by Walgreens is not a substitute for your own research concerning your healthcare, including Medicare.



Don't take chances with your health:

Fill ALL your prescriptions at one pharmacy.

At Walgreens we'll automatically screen for drug interactions and may be able to offer advice on money-saving alternatives.

For details, speak to our pharmacy staff.

Walgreens

There's a way to stay well.




NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

• Your Insurance Saved You: \$215.99

SUN 1:29PM
\$21.00
EXPRESS PAY

12/05/10
New



Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION

ABOUT YOUR MEDICATION.

Take advantage of these convenient services:

- **Auto Refills**
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.
- **Touch Tone Refills**
Save time by using our automated system for a refill. Just dial the number on your prescription label.

Walgreens

The Pharmacy America Trusts • Since 1901™


Visit us online at Walgreens.com

Thank you for choosing Walgreens!

Brunsting004490

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
12850 Memorial Drive
Houston, TX 77024
(713)722-7247

PATIENT	NELVA BRUNSTING	DOCTOR	A. JAIN, MD	DRUG DESCRIPTION	 PINK FRONT: m 32
BIRTH DATE	10/08/26	PATIENT ALLERGIES			
MEDICATION	METOPROLOL TARTRATE 50MG TABLETS				
QUANTITY	60				
DIRECTIONS	TAKE 1 TABLET BY MOUTH EVERY 12 HOURS				
	9 am 3 9 pm				

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

INGREDIENT NAME: METOPROLOL (me-TOE-proe-lole)

COMMON USES: This medicine is a beta-adrenergic blocking agent (beta-blocker) used to treat high blood pressure. It may be used alone or with other medicines. It is also used for the long-term treatment of chest pain (angina) and to reduce the risk of death due to heart problems in certain patients who have had a heart attack. It may also be used for other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: **WARNING:** Do not suddenly stop taking this medicine; sharp chest pain, irregular heartbeat, and sometimes heart attack may occur. The risk may be greater if you have certain types of heart disease. Your doctor should slowly lower your dose over several weeks if you need to stop taking it, even if you only take it for high blood pressure. Heart disease is common and you may not know you have it. If you have chest pain or other heart problems occur, contact your doctor right away. You may need to start taking this medicine again. Tell your doctor or dentist that you take this medicine before you receive any medical or dental care, emergency care, or surgery. Tell your doctor if you have a history of diabetes or take medicine to lower your blood sugar (eg, glyburide, insulin). This medicine may hide signs of low blood sugar such as fast heartbeat. Tell your doctor right away if you notice other signs of low blood sugar such as anxiety, chills, dizziness, drowsiness, fainting, headache, tremor, unusual sweating, vision changes, or weakness. This medicine should not usually be used by patients who have a history of certain lung or breathing problems (eg, asthma) or a certain type of adrenal gland tumor (pheochromocytoma). It may worsen these conditions. Tell your doctor if you have a history of these conditions; your doctor may need to adjust your dose or prescribe additional medicine to reduce the risk of side effects. Tell your doctor if you have a history of overactive thyroid. This medicine may hide symptoms of overactive thyroid (eg, fast heartbeat). Do not suddenly stop taking this medicine; suddenly stopping this medicine could worsen your condition. Your doctor should slowly lower your dose over several weeks if you need to stop taking it. Check with your doctor for more information. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking mifepristate. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking amiodarone, bupropion, certain HIV protease inhibitors (eg, ritonavir), certain selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), cimetidine, digoxin, diphendramine, disopyramide, flecainide, hormonal contraceptives (eg, birth control pills), hydralazine, hydroxychloroquine, ketanserin, mefloquine, phenothiazines (eg, thioridazine), propafenone, quinazoline (eg, aluzosin), quinidine, terbinafine, thiazines (eg, propylthiouracil, verapamil), barbiturates (eg, phenobarbital), indomethacin, phenylpropanolamine, bupivacaine, lidocaine, or clonidine. **DO NOT STOP** or change your medicine without your doctor or pharmacist approval. Inform your doctor of any other medical conditions including lung or breathing problems (eg, asthma, bronchitis, chronic obstructive pulmonary disease (COPD)), diabetes, low blood pressure, thyroid disease, or adrenal gland tumor (pheochromocytoma), allergies, pregnancy, or breastfeeding. Tell your doctor if you have a history of heart attack, slow or irregular heartbeat, heart failure, or other heart problems; chest pain or angina; blood vessel or circulation problems; liver disease; or if you will be having surgery or receiving anesthesia. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have a very slow heartbeat (eg, bradycardia), certain types of irregular heartbeat (eg, atrioventricular [AV] block, sick sinus syndrome), moderate to severe heart failure, very low systolic blood pressure (less than 100 mm Hg), or severe blood circulation problems. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for taking this medicine provided by your doctor. **TAKE THIS MEDICINE BY MOUTH WITH FOOD** or immediately following meal at the same time each day. **STORE THIS MEDICINE** at room temperature, between 59 and 86 degrees F (15 and 30 degrees C). Store away from heat, moisture, and light. Do not store in the bathroom. **IF YOU MISS A DOSE OF THIS MEDICINE,** take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

CAUTIONS: **DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine or to another beta-blocker (eg, propranolol). **IF YOU HAVE A HISTORY OF ANY SEVERE ALLERGIC REACTION,** talk with your doctor. You may be at risk for an even more severe allergic reaction if you come into contact with the substance that caused your allergy. Some medicines used to treat

severe allergies may also not work as well while you are using this medicine. **LAB TESTS,** including liver and kidney function, blood pressure, and complete blood cell counts, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to keep all doctor and lab appointments. **THIS MEDICINE MAY CAUSE DROWSINESS, DIZZINESS, OR LIGHTHEADEDNESS.** These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. Do not drive or perform other possibly unsafe tasks until you know how you react to it. This medicine may cause dizziness, lightheadedness, or fainting; alcohol, hot weather, exercise, or fever may increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of any of these effects. **PATIENTS WHO TAKE MEDICINE FOR HIGH BLOOD PRESSURE** often feel tired or run down for a few weeks after starting treatment. Be sure to take your medicine even if you may not feel "normal." Tell your doctor if you develop any new symptoms. **DO NOT SUDDENLY STOP TAKING THIS MEDICINE** without first consulting your doctor. If your doctor decides you should no longer use this medicine, you will need to stop this medicine gradually according to your doctor's instructions. **IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK YOUR BLOOD PRESSURE** and heart rate regularly, be sure to do so. **TELL YOUR DOCTOR OR DENTIST** that you take this medicine before you receive any medical or dental care, emergency care, or surgery. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescription or over-the-counter, check with your doctor or pharmacist. Do not take any medicines used to treat colds or congestion without first talking with your doctor or pharmacist. **FOR WOMEN:** If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. This medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby. **DIABETES PATIENTS:** this medicine may hide signs of low blood sugar, such as rapid heartbeat. Be sure to watch for other signs or low blood sugar. Low blood sugar may make you anxious, sweaty, weak, dizzy, drowsy, or faint. It may also make your vision change; give you a headache, chills, or tremors; or make you more hungry. Check blood sugar levels closely. Ask your doctor before you change the dose of your diabetes medicine.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while taking this medicine include constipation, diarrhea, dizziness, dry mouth/eyes, gas, headache, heartburn, lightheadedness, mild drowsiness, muscle aches, nausea, stomach pain, trouble sleeping, unusual tiredness or weakness, or vomiting. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience blue or unusually cold hands or feet; chest pain; fainting; hallucinations; mood or mental changes (eg, confusion, depression); pounding in the chest; severe dizziness; or lightheadedness; shortness of breath; slow or irregular heartbeat; swelling of the arms, hands, and feet; vision changes; wheezing; yellowing of the skin or eyes. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include chest pain, seizures, very slow heart beat, severe dizziness, fainting, and difficult or slowed breathing.

ADDITIONAL INFORMATION: **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** obtain refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

Do not flush unused medications or pour down a sink or drain. WICH 957918

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

NELVA BRUNSTING
13630 Pinerock, Houston, TX 770797517
(713)948-4391
RX # 1496588-03328 DATE: 12/05/10

METOPROLOL TARTRATE 50MG TABLETS
QTY: 60 3 REFILLS BEFORE 12/05/11
New NDC:00378-0032-10
Retail Price: \$14.99 Your Insurance Saved You: \$10.61 **\$ 4.38**

A. JAIN, MD PLAN: PAIDMPD
MFG:MYLAN GROUP#: CMD3896
TPL/RJW/RJW/ RJW CLAIM REF#: YRKMEWD

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024 Customer
PH: (713)722-7247 Receipt


NELVA BRUNSTING
13630 Pinerock, Houston, TX 770797517
(713)948-4391
RX # 1496588-03328 DATE: 12/05/10

METOPROLOL TARTRATE 50MG TABLETS
QTY: 60 3 REFILLS BEFORE 12/05/11
New NDC:00378-0032-10
Retail Price: \$14.99 Your Insurance Saved You: \$10.61 **\$ 4.38**

A. JAIN, MD PLAN: PAIDMPD
MFG:MYLAN GROUP#: CMD3896
TPL/RJW/RJW/ RJW CLAIM REF#: YRKMEWD

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024 Duplicate
PH: (713)722-7247 Receipt

Pharmacy use only

SUN	1:29PM	METOPROLOL TARTRATE 50MG TABLETS	QTY 60	 PINK FRONT: m 32
New		00378-0032-10	20 DRAM	
		CELL 138		

TPL/RJW/RJW/ RJW

Brunsting004481

Educate Yourself About Stroke

Stroke kills more women than breast cancer, cervical cancer, uterine cancer, and ovarian cancer combined¹

Be Prepared...

ACT F.A.S.T!

Stroke risk factors

FACE	Facial droop Uneven smile	High blood pressure High cholesterol Heart disease
ARM	Arm numbness Arm weakness	Diabetes Smoking
SPEECH	Slurred speech Difficulty speaking or understanding	Heavy alcohol use Physical inactivity and obesity
TIME	Call 911 and get to the hospital immediately.	Atrial fibrillation (irregular heartbeat) Family history of stroke

Check out National Stroke Association's
Virtual Health Fair, January 19, 2011 and beyond

- Register now and get comprehensive, accurate, and up-to-date education on and resources for stroke prevention and recovery
- Visit www.stroke.org/catalina, hosted by National Stroke Association

For more information on stroke,
visit www.getstrokeinfo.com

Reference: 1. Heron M, Hoyert DL, Murphy SL, Xu J, Kochanek KD, Tejada-Vera B. Deaths: final data for 2008. *Natl Vital Stat Rep*. 2009;57(14):1-134.

Brought to you by
Genentech
A Member of the Roche Group

©2010 Genentech USA, Inc. All rights reserved. AC1000107200 Printed in USA.

You're taking action
by managing the most important part of your life, your health.



Having your prescriptions refilled on time is a positive step towards managing your condition, but it's just one part of managing it. Eating healthy foods and getting proper exercise are also important factors for total health. If you don't already have a diet and exercise regimen, then talk to your doctor about the best plan for you.

You are taking **metoprolol** to help control your high blood pressure. Below are some key points to remember about **metoprolol**.

- Follow the directions on your prescription bottle carefully.
- Take your medicine exactly as your doctor prescribed.
- Continue to take it even if you feel well. Give your body time to adjust to the medicine.
- To help you remember, take it around the same time(s) every day.
- Check your blood pressure regularly.
- If you have any questions, ask one of our pharmacists on the pharmacy team or call your doctor. It's important for you to keep all scheduled appointments for lab testing.
- Talk to your doctor about how you feel and about any symptoms you have during the starting of this medicine.

The information above is brought to you as a courtesy from your pharmacist.

8
BR




SUN 1:29PM
\$4.38
EXPRESS PAY

NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

12/05/10
New

• Your Insurance Saved You: \$10.61



Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION

ABOUT YOUR MEDICATION.

Take advantage of these convenient services:

- **Touch Tone Refills**
Save time by using our automated system for a refill. Just dial the number on your prescription label.
- **Auto Refills**
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

Walgreens
The Pharmacy America Trusts • Since 1901™
Visit us online at Walgreens.com

Thank you for choosing Walgreens!

To stop receiving sponsored info at this pharmacy, call 888-336-5744 or visit remove-me.net Use code:2875677 339 0223746

Brunsting004482

AA Brunsting Financials 003942



OIRT: 2875 677 339 0223746

INFO: 0896 00378003210

3440

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
12850 Memorial Drive
Houston, TX 77024
(713)722-7247

PATIENT NELVA BRUNSTING BIRTH DATE 10/08/26 MEDICATION METOPROLOL TARTRATE 50MG TABLETS QUANTITY 60 DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY 12 HOURS	DOCTOR A. JAIN, MD PATIENT ALLERGIES  DRUG DESCRIPTION 
---	--

INGREDIENT NAME: METOPROLOL (me-TOE-proe-lole)

COMMON USES: This medicine is a beta-adrenergic blocking agent (beta-blocker) used to treat high blood pressure. It may be used alone or with other medicines. It is also used for the long-term treatment of chest pain (angina) and to reduce the risk of death due to heart problems in certain patients who have had a heart attack. It may also be used for other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: WARNING: Do not suddenly stop taking this medicine; sharp chest pain, irregular heartbeat, and sometimes heart attack may occur. The risk may be greater if you have certain types of heart disease. Your doctor should slowly lower your dose over several weeks if you need to stop taking it, even if you only take it for high blood pressure. Heart disease is common and you may not know you have it. Limit physical activity while you are lowering your dose. If new or worsened chest pain or other heart problems occur, contact your doctor right away. You may need to start taking this medicine again. Tell your doctor or dentist that you take this medicine before you receive any medical or dental care, emergency care, or surgery. Tell your doctor if you have a history of diabetes or take medicine to lower your blood sugar (eg, glyburide, insulin). This medicine may hide signs of low blood sugar such as fast heartbeat. Tell your doctor right away if you notice other signs of low blood sugar such as anxiety, chills, dizziness, drowsiness, fainting, headache, tremor, unusual sweating, vision changes, or weakness. This medicine should not usually be used by patients who have a history of certain lung or breathing problems (eg, asthma) or a certain type of adrenal gland tumor (pheochromocytoma). It may worsen these conditions. Tell your doctor if you have a history of these conditions; your doctor may need to adjust your dose or prescribe additional medicine to reduce the risk of side effects. Tell your doctor if you have a history of overactive thyroid. This medicine may hide symptoms of overactive thyroid (eg, fast heartbeat). Do not suddenly stop taking this medicine; suddenly stopping this medicine could worsen your condition. Your doctor should slowly lower your dose over several weeks if you need to stop taking it. Check with your doctor for more information about other medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking mifepridil. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking amiodarone, bupropion, certain HIV protease inhibitors (eg, ritonavir), certain selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), cimetidine, diphtheria (cramine, dipyramide, hecainide, certain HIV protease inhibitors (eg, birth control pills), hydroxyzine, hydroxychloroquine, ketanserin, mefloquine, phenothiazines (eg, thioridazine), propafenone, quinsalines (eg, alfuzosin), quinidine, terbinafine, thiamines (eg, prolylbouracil), verapamil, barbiturates (eg, phenobarbital), indomethacin, phenylpropranolamine, bupivacaine, lidocaine, or clonidine. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions including lung or breathing problems (eg, asthma, bronchitis, chronic obstructive pulmonary disease [COPD]), diabetes, low blood pressure, thyroid problems, or sick sinus syndrome, moderate to severe heart failure, very low systolic blood pressure (less than 100 mm Hg), or severe blood circulation problems. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for taking this medicine provided by your doctor. TAKE THIS MEDICINE BY MOUTH WITH FOOD or immediately following a meal at the same time each day. STORE THIS MEDICINE at room temperature, between 59 and 86 degrees F (15 and 30 degrees C). Store away from heat, moisture, and light. Do not store in the bathroom. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

CAUTIONS: DO NOT USE THIS MEDICINE if you are allergic to any ingredient in this medicine or to another beta-blocker (eg, propranolol). IF YOU HAVE A HISTORY OF ANY SEVERE ALLERGIC REACTION, talk with your doctor. You may be at risk for an even more severe allergic reaction if you come into contact with the substance that caused your allergy. Some medicines used to treat

severe allergies may also not work as well while you are using this medicine. **LAB TESTS**, including liver and kidney function, blood pressure, and complete blood cell counts, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to keep all doctor and lab appointments. **THIS MEDICINE MAY CAUSE DROWSINESS, DIZZINESS, OR LIGHTEADEDNESS.** These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. Do not drive or perform other possibly unsafe tasks until you know how you react to it. This medicine may cause dizziness, lightheadedness, or fainting; alcohol, hot weather, exercise, or fever may increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of any of these effects. **PATIENTS WHO TAKE MEDICINE FOR HIGH BLOOD PRESSURE** often feel tired or run down for a few weeks after starting treatment. Be sure to take your medicine even if you may not feel "normal." Tell your doctor if you develop any new symptoms. **DO NOT SUDDENLY STOP TAKING THIS MEDICINE** without first consulting your doctor. If your doctor decides you should no longer use this medicine, you will need to stop this medicine gradually according to your doctor's instructions. IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK YOUR BLOOD PRESSURE and heart rate regularly, be sure to do so. **TELL YOUR DOCTOR OR DENTIST** that you take this medicine before you receive any medical or dental care, emergency care, or surgery. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES**, either prescription or over-the-counter, check with your doctor or pharmacist. Do not take any medicines used to treat colds or congestion without first talking with your doctor or pharmacist. **FOR WOMEN:** If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. This medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby. **DIABETES PATIENTS:** this medicine may hide signs of low blood sugar, such as rapid heartbeat. Be sure to watch for other signs of low blood sugar. Low blood sugar may make you anxious, sweaty, weak, dizzy, drowsy, or faint. It may also make your vision change; give you a headache, chills, or tremors; or make you more hungry. Check blood sugar levels closely. Ask your doctor before you change the dose of your diabetes medicine.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while taking this medicine include constipation, diarrhea, dizziness, dry mouth/eyes, gas, headache, heartburn, lightheadedness, mild drowsiness, muscle aches, nausea, stomach pain, trouble sleeping, unusual tiredness or weakness, or vomiting. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience blue or unusually cold hands or feet; chest pain; fainting; hallucinations; mood or mental changes (eg, confusion, depression); pounding in the chest; severe dizziness or lightheadedness; shortness of breath; slow or irregular heartbeat; swelling of the arms, hands, and feet; vision changes; wheezing; yellowing of the skin or eyes. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include chest pain, seizures, very slow heart beat, severe dizziness, fainting, and difficult or slowed breathing.

ADDITIONAL INFORMATION: DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

NELVA BRUNSTING

13630 Pinerock, Houston, TX 770797517
7713464-4391

RX # 1496588-03328

DATE: 01/01/11

METOPROLOL TARTRATE 50MG TABLETS

QTY: 60 2 REFILLS BEFORE 12/05/11

Refill NDC:00378-0032-10

Retail Price: \$14.99 Your Insurance Saved You: \$10.61

\$ 4.38

A. JAIN, MD
MFG:MYLAN
XXX/JIC/JIC /R/JW

PLAN: PAIDMD
GROUP# CMD3896
CLAIM REF# YXEP3F1

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
PH: (713)722-7247

Customer
Receipt

NELVA BRUNSTING

13630 Pinerock, Houston, TX 770797517
7713464-4391

RX # 1496588-03328

DATE: 01/01/11

METOPROLOL TARTRATE 50MG TABLETS

QTY: 60 2 REFILLS BEFORE 12/05/11

Refill NDC:00378-0032-10

Retail Price: \$14.99 Your Insurance Saved You: \$10.61

\$ 4.38

A. JAIN, MD
MFG:MYLAN
XXX/JIC/JIC /R/JW

PLAN: PAIDMD
GROUP# CMD3896
CLAIM REF# YXEP3F1

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
PH: (713)722-7247

Duplicate
Receipt

Pharmacy use only

METOPROLOL TARTRATE 50MG TABLETS
SUN 12:00PM 00378-0032-10
Refill CELL 138

QTY 60
20 DRAM



PINK
FRONT: m 32

XXX/JIC/JIC /R/JW

Brunsting004479

AABrunsting.Financials003944

Educate Yourself About Stroke

Stroke kills more women than breast cancer, cervical cancer, uterine cancer, and ovarian cancer combined¹

Be Prepared...

ACT F.A.S.T!

Stroke risk factors

F FACE	Facial droop Uneven smile	High blood pressure High cholesterol Heart disease
A ARM	Arm numbness Arm weakness	Diabetes Smoking
S SPEECH	Slurred speech Difficulty speaking or understanding	Heavy alcohol use Physical inactivity and obesity
T TIME	Call 911 and get to the hospital immediately.	Atrial fibrillation (irregular heartbeat) Family history of stroke

Check out National Stroke Association's Virtual Health Fair, January 19, 2011 and beyond

- Register now and get comprehensive, accurate, and up-to-date education on and resources for stroke prevention and recovery
- Visit www.stroke.org/catalina, hosted by National Stroke Association

For more information on stroke,
visit www.getstrokeinfo.com

Reference: 1. Heron M, Hoyert DL, Murphy SL, Xu J, Kochanek KD, Tejada-Vera B. Deaths: final data for 2008. *Natl Vital Stat Rep*. 2008;57(14):1-134.

Brought to you by **Genentech**
A Member of the Roche Group

©2010 Genentech USA, Inc. All rights reserved. AC10800107200 Printed in USA.

You're taking action

by managing the most important part of your life, your health.



Having your prescriptions refilled on time is a positive step towards managing your condition, but it's just one part of managing it. Eating healthy foods and getting proper exercise are also important factors for total health. If you don't already have a diet and exercise regimen, then talk to your doctor about the best plan for you.

You are taking **metoprolol** to help control your high blood pressure. Below are some key points to remember about **metoprolol**.

- Follow the directions on your prescription bottle carefully.
- Take your medicine exactly as your doctor prescribed.
- Continue to take it even if you feel well. Give your body time to adjust to the medicine.
- To help you remember, take it around the same time(s) every day.
- Check your blood pressure regularly.
- If you have any questions, ask one of our pharmacists on the pharmacy team or call your doctor. It's important for you to keep all scheduled appointments for lab testing.
- Talk to your doctor about how you feel and about any symptoms you have during the starting of this medicine.

The information above is brought to you as a courtesy from your pharmacist.

8
BR




SUN 12:00PM
\$4.38
EXPRESS PAY

NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

01/01/11
Refill

- Store TT 1 of 5
- Your Insurance Saved You: \$10.61



Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION

ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- **Auto Refills**
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

Walgreens

The Pharmacy America Trusts • Since 1901*

Visit us online at Walgreens.com

Thank you for choosing Walgreens!

To stop receiving sponsored info at this pharmacy, call 888-336-5744 or visit remove-me.net Use code:2875677 002 0240051

Brunsting004480

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
12850 Memorial Drive
Houston, TX: 77024
(713)722-7247

<p>PATIENT NELVA BRUNSTING BIRTH DATE 10/08/26 MEDICATION ETHAMBUTOL 400MG TABLETS QUANTITY 90 DIRECTIONS TAKE 3 TABLETS BY MOUTH EVERY DAY</p>	<p>DOCTOR A. JAIN, MD PATIENT ALLERGIES</p>	<p>DRUG DESCRIPTION</p> <div style="border: 1px solid black; width: 40px; height: 40px; margin: 0 auto; display: flex; align-items: center; justify-content: center;"> ○ </div> <p style="text-align: center;">WHITE FRONT: L U BACK: C32</p>
<p>INGREDIENT NAME: ETHAMBUTOL (e-THAM-byoo-tole)</p> <p>COMMON USES: This medicine is an antibacterial used to treat tuberculosis (TB).</p> <p>BEFORE USING THIS MEDICINE: INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. Inform your doctor of any other medical conditions, allergies, pregnancy, or breast-feeding.</p> <p>HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. STORE THIS MEDICINE at room temperature in a tightly-closed container, away from heat and light. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.</p> <p>CAUTIONS: DO NOT STOP USING THIS MEDICINE without first checking with your doctor. THIS MEDICINE MAY CAUSE dizziness. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. HAVE REGULAR EYE EXAMINATIONS while you are taking this medicine even if you do not notice changes in your vision. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist.</p> <p>POSSIBLE SIDE EFFECTS: SIDE EFFECTS, that may go away during</p>		
<p>treatment, include nausea, headache, or dizziness. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience changes in vision, vomiting, skin rash, itching, fever, confusion, or numbness or tingling of extremities. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.</p> <p>OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately.</p> <p>ADDITIONAL INFORMATION: If your symptoms do not improve within 2 to 3 weeks, or if they become worse, check with your doctor. DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out.</p>		
<p>KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.</p>		

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain. WIC# 957918

NELVA BRUNSTING
13630 Finerock, Houston, TX 770797517
(713)464-4381
RX # 1494792-03328 DATE: 01/01/11

ETHAMBUTOL 400MG TABLETS
QTY: 90 2 REFILLS BEFORE 11/29/11
Refill NDC: 68180-0281-01
Retail Price: \$153.59 Your Insurance Saved You: \$148.59 **\$ 5.00**

A. JAIN, MD PLAN: PAIDMPD
MFG: LUPIN GROUP# CMD3896
XXX/JIC/JIC/ RJW CLAIM REF# SQA93NQ

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024 **Customer Receipt**
PH: (713)722-7247

NELVA BRUNSTING
13630 Finerock, Houston, TX 770797517
(713)464-4381
RX # 1494792-03328 DATE: 01/01/11

ETHAMBUTOL 400MG TABLETS
QTY: 90 2 REFILLS BEFORE 11/29/11
Refill NDC: 68180-0281-01
Retail Price: \$153.59 Your Insurance Saved You: \$148.59 **\$ 5.00**

A. JAIN, MD PLAN: PAIDMPD
MFG: LUPIN GROUP# CMD3896
XXX/JIC/JIC/ RJW CLAIM REF# SQA93NQ

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024 **Duplicate Receipt**
PH: (713)722-7247

<p>Pharmacy use only</p> <p>SUN 12:00PM Refill</p>	<p>ETHAMBUTOL 400MG TABLETS 68180-0281-01 ALPHA</p>	<p>QTY 90</p>	<div style="border: 1px solid black; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center;"> ○ </div> <p style="text-align: center;">WHITE FRONT: L U BACK: C32</p> <p style="text-align: center; font-size: 0.8em;">XXX/JIC/JIC/ RJW</p>
---	---	---------------	--

Brunsting004473

Get a **FREE** one-on-one
Medicare Part D
review session!

Your pharmacist will:

- Review your medications and look for ways to help save you money
- Talk about other things you can do to stay well
- Check for potential drug interactions



Make your appointment today!


Walgreens

There's a way to stay well.

Information provided by Walgreens is not a substitute for your own research concerning your healthcare, including Medicare.

Fever at 3 a.m.?

We have the most 24-hour pharmacies in America, so you get the help you need, when you need it.



For the 24-hour Walgreens nearest you, call 1-800-WALGREENS or visit Walgreens.com/findastore.

Another great service from

Walgreens

The Pharmacy America Trusts Since 1901

8
BR



NELVA BRUNSTING

13630 Pinerock
Houston, TX 770797517
(713)464-4391

- Store TT 2 of 5
- Your Insurance Saved You: \$148.59

SUN 12:00PM
\$5.00
EXPRESS PAY

01/01/11
Refill



**Personal
Prescription
Information**

LOOK INSIDE FOR IMPORTANT INFORMATION

ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- Auto Refills
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

Walgreens

The Pharmacy America Trusts • Since 1901™


Visit us online at Walgreens.com

Thank you for choosing Walgreens!

Brunsting004474

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
 12850 Memorial Drive
 Houston, TX 77024
 (713)722-7247

PATIENT	NELVA BRUNSTING	DOCTOR	A. JAIN, MD	DRUG DESCRIPTION
BIRTH DATE	10/08/26	PATIENT ALLERGIES		
MEDICATION	AZITHROMYCIN 250MG TABLETS			PINK
QUANTITY	30			FRONT: 93
DIRECTIONS	TAKE 1 TABLET BY MOUTH EVERY DAY			BACK: 7146

INGREDIENT NAME: AZITHROMYCIN
 (ay-ZITH-roe-MYE-sin)

COMMON USES: This medicine is a macrolide antibiotic used to treat bacterial infections.

BEFORE USING THIS MEDICINE: Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking dofetilide, nilotinib, propafenone, pimozide, or tetrabenazine. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking antiarrhythmics (eg, disopyramide), anticoagulants (eg, warfarin), arsenic, astemizole, carbamazepine, cisapride, digoxin, domperidone, maprotiline, methadone, nelfinavir, cyclosporine, ergot alkaloids (eg, ergotamine), paliperidone, phenytoin, quinolones (eg, levofloxacin), rifampin, terfenadine, theophylline, triazolam, tyrosine kinase inhibitors (eg, dasatinib), or medicines that may affect your heartbeat. Ask your doctor if you are unsure if any of the medicines you are taking may affect your heartbeat. Inform your doctor of any other medical conditions including irregular heartbeat, kidney problems, liver problems, myasthenia gravis, allergies, pregnancy or breastfeeding. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine may be taken on an empty stomach or with food. DO NOT TAKE THIS MEDICINE within 1 hour before or 2 hours after aluminum- or magnesium-containing antacids. STORE THIS MEDICINE at room temperature, away from heat and light. TO CLEAR UP YOUR INFECTION COMPLETELY, continue taking this medicine for the full course of treatment even if you feel better in a few days. Do not miss any doses. Taking this medicine at the same time each day will make it easier to remember. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. If you miss a dose, do not take 2 doses at once.

CAUTIONS: DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it, to other macrolide antibiotics (such as erythromycin), to ketolide antibiotics (such as telithromycin), or if you are allergic to any ingredient in this product. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to erythromycin or any macrolide or ketolide antibiotic. A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine, contact your doctor or pharmacist. IF YOU EXPERIENCE difficulty breathing; tightness of chest; swelling of eyelids, face, or lips; or if you develop a rash or hives, tell your doctor immediately. Do not take any more of this medicine unless your doctor tells you to do so. This medicine may cause drowsiness, dizziness, or lightheadedness. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. THIS

MEDICINE MAY CAUSE increased sensitivity to the sun. Avoid exposure to the sun, sunlamps, or tanning booths until you know how you react to this medicine. Use a sunscreen or protective clothing if you must be outside for a prolonged period. MILD DIARRHEA IS COMMON with antibiotic use. However, a more serious form of diarrhea (pseudomembranous colitis) may rarely occur. This may develop while you use the antibiotic or within several months after you stop using it. Contact your doctor right away if stomach pain or cramps, severe diarrhea, or bloody stools occur. Do not treat diarrhea without first checking with your doctor. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. FOR WOMEN: IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. IT IS UNKNOWN IF THIS MEDICINE IS EXCRETED in breast milk. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS, that may go away during treatment, include mild diarrhea, headache, nausea, or stomach pain or upset. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience vomiting, changes in hearing, or ringing in the ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience chest pain; swelling of your hands, legs, face, lips, eyes, throat, or tongue; difficulty swallowing, speaking, or breathing; eye or vision problems; fainting; hearing loss; irregular heartbeat; muscle weakness; reddened, blistered, or swollen skin; seizures; severe stomach pain, cramps, or diarrhea; bloody stools; unusual vaginal itching, odor, or discharge; or yellowing of the eyes or skin. An allergic reaction to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, dizziness, trouble breathing, or unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include nausea, vomiting, stomach upset, and diarrhea.

ADDITIONAL INFORMATION: DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

NELVA BRUNSTING
 13630 Pinerock, Houston, TX 770797517
 (713)464-4391
RX # 1494789-03328 DATE: 01/01/11

AZITHROMYCIN 250MG TABLETS
 QTY: 30 2 REFILLS BEFORE 11/29/11
 Refill NDC: 00093-7146-56
 Retail Price: \$195.79 Your Insurance Saved You: \$190.79

\$ 5.00

A. JAIN, MD PLAN: PAIDMPO
 MFG: TEVA GROUP# CMD3886
 XXX/KHN/KHN/KHN/RJW CLAIM REF# SQA93N1

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247

Customer Receipt

NELVA BRUNSTING
 13630 Pinerock, Houston, TX 770797517
 (713)464-4391
RX # 1494789-03328 DATE: 01/01/11

AZITHROMYCIN 250MG TABLETS
 QTY: 30 2 REFILLS BEFORE 11/29/11
 Refill NDC: 00093-7146-56
 Retail Price: \$195.79 Your Insurance Saved You: \$190.79

\$ 5.00

A. JAIN, MD PLAN: PAIDMPO
 MFG: TEVA GROUP# CMD3886
 XXX/KHN/KHN/KHN/RJW CLAIM REF# SQA93N1

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

SUN 12:00PM AZITHROMYCIN 250MG TABLETS
 Refill 00093-7146-56
 ALPHA

QTY 30
 10 DRAM



PINK
 FRONT: 93
 BACK: 7146

XXX/KHN/KHN/KHN/RJW

Brunsting004475

AABrunsting.Financials003948



Are you getting the most from your Medicare Part D plan?

We can help you find out with a **FREE one-on-one plan review session** with your pharmacist. We'll review your medications, look for ways to help save you money and more.

Make your appointment today!

Walgreens

There's a way to stay well.

Information provided by Walgreens is not a substitute for your own research concerning your healthcare, including Medicare.

49776

AARP Members

Get special offers each week from Walgreens!*

Just show your AARP card at time of purchase to receive exclusive offers on everyday products - available only to AARP members.

AARP | Walgreens
Smarter. Stronger. Healthier.

*AARP card must be presented at time of purchase. For complete program terms and conditions, visit Walgreens.com/aarpcard. AARP receives a royalty for the use of its intellectual property. Amounts paid are used for the general purposes of AARP and its members.

DPF: 2875 677 002 0740067

INFO: 0900 00093714656

5434



NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

- Store TT 3 of 5
- Your Insurance Saved You: \$190.79

SUN 12:00PM
\$5.00
EXPRESS PAY

01/01/11
Refill

Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION

ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- Auto Refills
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

Walgreens

The Pharmacy America Trusts • Since 1901™

Visit us online at Walgreens.com

Thank you for choosing Walgreens!


Brunsting004476

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
12850 Memorial Drive
Houston, TX 77024
(713)722-7247

<p>PATIENT NELVA BRUNSTING BIRTH DATE 10/08/26 MEDICATION RIFAMPIN 300MG CAPSULES QUANTITY 60 DIRECTIONS TAKE 2 CAPSULES BY MOUTH EVERY DAY</p>	<p>DOCTOR A. JAIN, MD PATIENT ALLERGIES</p>
--	--

DRUG DESCRIPTION



DARK REDDISH-BROWN
FRONT: LANNETT
BACK: 1315

INGREDIENT NAME: RIFAMPIN (rif-AM-pin)

COMMON USES: This medicine is a rifamycin antibiotic used to treat tuberculosis (TB), as well as to treat those who have been exposed to meningitis-causing bacteria before they become sick. This drug does not treat acute meningitis. It may also be used to treat other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking HIV protease inhibitors, pyrazinamide, or birth control pills. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking anticoagulants, cyclosporine, digitoxin, isoniazid, itraconazole, macrolide antibiotics, mexiletine, nevirapine, quinidine, progestins, theophylline, tocinamide, verapamil, or medicine for anxiety, sleep, or seizures. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, allergies, pregnancy, or breast-feeding. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. Take this medicine on an empty stomach at least 1 hour before or 2 hours after eating. **THIS MEDICINE MAY BE TAKEN WITH FOOD** if it upsets your stomach. **STORE THIS MEDICINE** at room temperature in a tightly-closed container, away from heat and light. **TO CLEAR UP YOUR INFECTION COMPLETELY**, continue taking this medicine for the full course of treatment. Do not miss any doses. **IF YOU MISS A DOSE OF THIS MEDICINE**, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

CAUTIONS: IT MAY TAKE SEVERAL WEEKS for this medicine to work. Do not stop using this medicine without checking with your doctor. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. **THIS MEDICINE MAY CAUSE** urine, feces, saliva, sweat, and tears to turn orange or red. **THIS MEDICINE MAY PERMANENTLY STAIN** soft contact lenses. **KEEP ALL DOCTOR AND LABORATORY**

APPOINTMENTS while you are using this medicine. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE**, either prescription or over-the-counter, check with your doctor or pharmacist. **USING THIS MEDICINE** while you are taking birth control pills may decrease the effectiveness of your birth control pills. To prevent pregnancy, use an additional form of birth control. **FOR WOMEN: IF YOU PLAN ON BECOMING PREGNANT**, discuss with your doctor the benefits and risks of using this medicine during pregnancy.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS, that may go away during treatment, include stomach upset, heartburn, loss of appetite, nausea, gas, abdominal cramping, headache, drowsiness, dizziness, menstrual changes, joint pain, or leg cramps. If they continue or are bothersome, check with your doctor. **CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE** if you experience vomiting, diarrhea, fever, chills, changes in vision, or confusion. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience severe stomach pain, dark urine, or yellowing of eyes or skin. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include swelling of face or around eyes, itching over the entire body, orange or red discoloration of skin or eyes, nausea, vomiting, drowsiness, and loss of consciousness.

ADDITIONAL INFORMATION: **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME**, obtain refills before your supply runs out.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain. WIC# 957918

NELVA BRUNSTING
 13630 Pinrock, Houston, TX 770797517
 (713)464-4391
RX # 1494790-03328 DATE: 01/01/11

RIFAMPIN 300MG CAPSULES
 QTY: 60 2 REFILLS BEFORE 11/29/11
 Refill NDC: 00527-1315-30
 Retail Price: \$113.89 Your Insurance Saved You: \$108.89 \$ 5.00

A. JAIN, MD PLAN: PAIDMPD
 MFGLANNETT GROUP# CMD3898
 XXX/KHN/KHN/KHN/RJW CLAIM REF# ONA1TWP

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247 Customer Receipt

NELVA BRUNSTING
 13630 Pinrock, Houston, TX 770797517
 (713)464-4391
RX # 1494790-03328 DATE: 01/01/11


RIFAMPIN 300MG CAPSULES
 QTY: 60 2 REFILLS BEFORE 11/29/11
 Refill NDC: 00527-1315-30
 Retail Price: \$113.89 Your Insurance Saved You: \$108.89 \$ 5.00

A. JAIN, MD PLAN: PAIDMPD
 MFGLANNETT GROUP# CMD3898
 XXX/KHN/KHN/KHN/RJW CLAIM REF# ONA1TWP

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247 Duplicate Receipt

Pharmacy use only

<p>SUN 12:00PM Refill</p>	<p>RIFAMPIN 300MG CAPSULES 00527-1315-30 ALPHA</p>	<p>QTY 60 20 DRAM</p>
-------------------------------	--	---------------------------

 **DARK REDDISH-BROWN**
FRONT: LANNETT
BACK: 1315

XXX/KHN/KHN/KHN/RJW

Brunsting004477



Are you getting the most from your Medicare Part D plan?

We can help you find out with a **FREE one-on-one plan review session** with your pharmacist. We'll review your medications, look for ways to help save you money and more.

Make your appointment today!


Walgreens
There's a way to stay well.

Information provided by Walgreens is not a substitute for your own research concerning your healthcare, including Medicare.

61489

Prescription Transfers Easy as 1-2-3

1. Gather your prescription bottles.
2. Call or visit your local Walgreens.
3. We'll take care of the rest!



Avoid potentially harmful drug interactions by filling all your prescriptions at one pharmacy. By choosing Walgreens, our pharmacists can screen your prescriptions and warn you of potential interactions.

**For the Walgreens nearest you,
call 1-800-WALGREENS
or visit Walgreens.com/findastore.**

Another great service from
Walgreens
The Pharmacy America Trusts • Since 1901®

OPT: 2875 677 002 024080

INFO: 0990 0052731350

13689

8
BR




NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

- Store TT 4 of 5
- Your Insurance Saved You: \$108.89

SUN 12:00PM
\$5.00
EXPRESS PAY

01/01/11
Refill



Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION

ABOUT YOUR MEDICATION.
Take advantage of this convenient service:

- **Auto Refills**
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

Walgreens
The Pharmacy America Trusts • Since 1901®
Visit us online at Walgreens.com

Thank you for choosing Walgreens!

Brunsting004478

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
12850 Memorial Drive
Houston, TX 77024
(713)722-7247

PATIENT NELVA BRUNSTING
BIRTH DATE 10/08/26
MEDICATION SPIRIVA CAPS 30'S & HANDIHALER
QUANTITY 30
DIRECTIONS INHALE CONTENTS OF ONE CAPSULE ONCE DAILY USING HANDIHALER

DOCTOR A. JAIN, MD
PATIENT ALLERGIES

DRUG DESCRIPTION

INGREDIENT NAME: TIOTROPIUM BROMIDE (TYE-oh-TROE-pee-um BROE-mide)

COMMON USES: This medicine is an anticholinergic agent used to treat the symptoms of chronic obstructive pulmonary disease, including chronic bronchitis and emphysema. It may also be used to treat other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking another anticholinergic (eg, ipratropium). Ask your doctor if you are unsure if any of your medicines are anticholinergics. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medicine you are taking if you have trouble urinating; an enlarged prostate; bladder blockage; glaucoma; kidney problems; allergies (including milk proteins); pregnancy; or breast-feeding. Tell your doctor if you are having an asthma attack or increased difficulty breathing. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. An additional patient information leaflet is available with this medicine. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have before using this medicine. **DO NOT SWALLOW THE CAPSULES.** THE CAPSULES ARE USED WITH A SPECIAL INHALER. **DO NOT** remove a capsule from the packaging until you are ready to use it. To remove a capsule from the packaging, carefully peel the foil back to expose the capsule, then tip the capsule out of the blister. **DO NOT** cut the foil or use sharp objects to remove the capsule from the blister. If a second capsule is exposed to the air when you are removing a capsule or use, it must be discarded. **DO NOT** save the capsule for later. Place the capsule in the special inhaler device immediately. The device will puncture the capsule so that the medicine inside can be inhaled into the lungs through the mouthpiece. **EXHALE** slowly and deeply. **DO NOT** breathe into the mouthpiece of the inhaler. Position the inhaler mouthpiece between your lips and try to rest your tongue flat. Keep your head upright. Do not block the air vents on the inhaler. **TAKE A SLOW, DEEP BREATH.** You should hear or feel the capsule vibrate inside the inhaler. Hold your breath as long as it feels comfortable, then exhale slowly through pursed lips. Breathe out completely. To be sure all of the medicine has been inhaled from the capsule, inhale from the mouthpiece a second time following the same process. **ASK YOUR DOCTOR OR PHARMACIST** if you are unclear on how to use this device or inhale the medicine. **CLEAN THE INHALER DEVICE** once a month according to the instructions in the patient leaflet. Each inhaler device may be used for up to 1 year and then should be replaced. **STORE THIS MEDICINE** at room temperature at 77 degrees F (25 degrees C), away from heat, moisture, and light. Brief storage between 59 and 86 degrees F (15 and 30 degrees C) is permitted. Do not store in the bathroom. This medicine will work best if used regularly. Taking your dose at the same time each day will help you to remember. **IF YOU MISS A DOSE** OF THIS MEDICINE, use it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **DO NOT** use 2 doses at once. **DO NOT** use this medicine more often than 1 time every 24 hours.

CAUTIONS: **DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine or to atropine or related medicines (eg, ipratropium). **THIS MEDICINE WILL NOT STOP AN ASTHMA ATTACK** once one has started. **IF YOU ARE ALSO USING A RESCUE INHALER** (eg, albuterol), be sure to always carry the bronchodilator inhaler with you to use during asthma attacks. **DO NOT EXCEED THE RECOMMENDED DOSE** without checking with your doctor. **DO NOT STOP USING THIS MEDICINE** without first checking with your doctor. This medicine may cause dizziness or blurred vision. These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. **AVOID GETTING THIS MEDICINE IN YOUR EYES.** If you get the medicine in your eyes and eye pain, blurred vision, or other vision changes occur, contact your doctor immediately. **THIS MEDICINE MAY SOMETIMES CAUSE SEVERE BREATHING PROBLEMS** right after you use a dose. If this happens, use your short-acting bronchodilator inhaler. Contact your doctor or seek other medical care at once. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE**, either prescription or over-the-counter, check with your doctor or pharmacist. **USE THIS MEDICINE WITH CAUTION IN THE ELDERLY;** they may be more sensitive to its effects, especially constipation and urinary tract infections. **FOR WOMEN:** IF YOU

PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. **IT IS NOT KNOWN IF THIS MEDICINE IS FOUND** in breast milk. **IF YOU ARE OR WILL BE BREAST-FEEDING** while you are using the medicine, check with your doctor or pharmacist to discuss the risks to your baby.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while using this medicine include dry mouth, constipation, sinus inflammation, throat irritation, stomach pain, vomiting, blurred vision, or mild nosebleed. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience irritation, pain, or white patches in your mouth or on your tongue; mouth sores; severe or persistent nosebleeds; severe or persistent constipation; difficult or painful urination; eye pain or discomfort; vision changes (eg, halos, colored images); chest pain; fast or irregular heartbeat; difficulty swallowing; new or worsened breathing problems; or wheezing. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include mental changes; severe constipation; stomach pain; or tremors.

ADDITIONAL INFORMATION: **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** obtain refills before your supply runs out.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

NELVA BRUNSTING

13630 Pineroct, Houston, TX 770797517
(713)464-4391

RX # 1496586-03328

DATE: 01/01/11

SPIRIVA CAPS 30'S & HANDIHALER

QTY: 30 2 REFILLS BEFORE 12/05/11

Refill NDC: 00597-0075-41

Retail Price: \$260.99 Your Insurance Saved You: \$44.45

\$ 216.54

A. JAIN, MD
MFG: BOEHRINGER
XXX/JIC/JIC /RJW

PLAN: PATIDWPD
GROUP# CMD3896
CLAIM REF# MRWHDC

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
PH: (713)722-7247

Customer Receipt

NELVA BRUNSTING

13630 Pineroct, Houston, TX 770797517
(713)464-4391

RX # 1496586-03328

DATE: 01/01/11

SPIRIVA CAPS 30'S & HANDIHALER

QTY: 30 2 REFILLS BEFORE 12/05/11

Refill NDC: 00597-0075-41

Retail Price: \$260.99 Your Insurance Saved You: \$44.45

\$ 216.54

A. JAIN, MD
MFG: BOEHRINGER
XXX/JIC/JIC /RJW

PLAN: PATIDWPD
GROUP# CMD3896
CLAIM REF# MRWHDC

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

SUN 12:00PM
Refill

SPIRIVA CAPS 30'S & HANDIHALER
00597-0075-41
ALPHA

QTY 30

XXX/JIC/JIC /RJW

Brunsting004469

AABrunsting.Financials003952

Get a **FREE** one-on-one
Medicare Part D
review session!

Your pharmacist will:

- Review your medications and look for ways to help save you money
- Talk about other things you can do to stay well
- Check for potential drug interactions



Make your appointment today!

Walgreens

There's a way to stay well.

Information provided by Walgreens is not a substitute for your own research concerning your healthcare, including Medicare.

01/01/11

**Don't take
chances with
your health.**

Fill all your prescriptions
at one pharmacy

Walgreens automatically screens for drug interactions, including any over-the-counter medicines or supplements you tell us about.



**Play it safe. Fill all your
prescriptions at Walgreens.**

OFF: 2875.677.002.0240098

INFC: 0990.0059707541

3/2011

8
BR




NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

- Store TT 5 of 5
- Your Insurance Saved You: \$44.45

SUN 12:00PM
\$216.54
EXPRESS PAY

01/01/11
Refill



Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION

ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- **Auto Refills**
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

Walgreens

The Pharmacy America Trusts • Since 1901™

Visit us online at Walgreens.com

Thank you for choosing Walgreens!

Brunsting004470

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
12850 Memorial Drive
Houston, TX 77024
(713)722-7247

PATIENT	NELVA BRUNSTING	DOCTOR	A. JAIN, MD	DRUG DESCRIPTION	
BIRTH DATE	10/08/26	PATIENT ALLERGIES		LIQUID	
MEDICATION	BROVANA 15MCG/2ML INH SOL 30X2ML				
QUANTITY	120				
DIRECTIONS	INHALE 1 VIAL VIA NEBULIZER TWOICE DAILY				

INGREDIENT NAME: ARFORMOTEROL (ar-for-MOE-ter-ol)

COMMON USES: This medicine is a long-acting beta-agonist bronchodilator used for long-term treatment of chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema. It may also be used to treat other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: WARNING: LONG-ACTING BETA-AGONISTS SUCH AS THIS MEDICINE HAVE BEEN RARELY ASSOCIATED WITH AN INCREASED RISK OF ASTHMA-RELATED DEATH. Long-acting beta-agonists should not be used in asthma patients without another long-term asthma-control medicine (eg, inhaled corticosteroids). This medicine has not been approved to treat asthma. Safety and effectiveness of this medicine in patients with asthma have not been confirmed. **SOME MEDICINES MAY INTERACT** with this medicine. **DO NOT TAKE THIS MEDICINE** if you are using another medicine that has a long-acting beta-agonist (eg, salmeterol) in it. **TELL YOUR HEALTH CARE PROVIDER** if you are taking any other medicine, especially any of the following: corticosteroids (eg, prednisone), diuretics (eg, furosemide, hydrochlorothiazide), xanthines (eg, theophylline), catechol-O-methyltransferase (COMT) inhibitors (eg, entacapone), monoamine oxidase inhibitors (MAOI) (eg, phenelzine), or beta-blockers (eg, propranolol). This may not be a complete list of all interactions that may occur. Tell your doctor if you have taken any of these medicines may interact with other medicines you take. **DO NOT START OR STOP any medicine without doctor or pharmacist approval.** Inform your doctor of any other medical conditions, including a history of other breathing problems (eg, asthma), diabetes, heart problems (eg, fast or irregular heartbeat, heart blood vessel problems), liver problems, high blood pressure, low blood potassium levels, seizures, an overactive thyroid, or you have high blood or urine ketone levels, allergies, pregnancy, or breast-feeding. Tell your doctor if you have recently been to an emergency room for breathing problems; have a history of frequent hospitalizations for breathing problems; have ever had life-threatening breathing problems; or have had an unusual reaction to a sympathomimetic medicine (eg, albuterol, pseudoephedrine), such as fast or irregular heartbeat, overexcitement, or severe trouble sleeping. Tell your doctor if you have taken MAOI (eg, phenelzine) or a tricyclic antidepressant (eg, amitriptyline) within the last 14 days. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you are having severe breathing problems (eg, sudden, severe onset or worsening of COPD symptoms such as chest tightness, cough, shortness of breath, wheezing), you have asthma and you are not currently using a long-term asthma-control medicine (eg, inhaled corticosteroids), or if you have asthma that is already well controlled with the use of a long-term asthma-control medicine. **THIS MEDICINE SHOULD NOT BE USED IN CHILDREN;** safety and effectiveness in children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine comes with a MEDICATION GUIDE approved by the U.S. Food and Drug Administration. Read it carefully each time you refill this medicine. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. **THIS MEDICINE SHOULD ONLY BE INHALED USING A NEBULIZER.** Do not inhale or swallow it. A health care provider will teach you how to use the nebulizer. Be sure you know what type of nebulizer to use with this medicine, and how to use it. Follow the procedure you are taught when you use a dose. Contact your health care provider if you have any questions. **STORE THIS MEDICINE** in the original foil pouch in a dry place. Do not remove from the foil pouch of the vial until right before use. **IF THE MEDICINE CONTAINS PARTICLES,** is cloudy or discolored, or if the vial is cracked or damaged in any way, do not use it. **DO NOT MIX THIS MEDICINE WITH OTHER MEDICINES** in your nebulizer machine. **TO USE THIS MEDICINE,** twist open the top of the vial and pour the entire contents into the nebulizer reservoir. **CONNECT THE NEBULIZER RESERVOIR** to the mouthpiece or face mask. **CONNECT THE NEBULIZER** to the compressor. Sit in a comfortable, upright position. **PLACE THE MOUTHPIECE** in your mouth (or put on the face mask) and turn on the nebulizer. **BREATHE AS CALMLY,** deeply, and evenly as possible. **INHALE** the medicine in the nebulizer chamber (about 5 to 10 minutes). **CLEAN THE NEBULIZER** according to the instructions. Failure to properly clean the nebulizer could lead to bacteria entering the medicine. This may lead to an infection. To avoid bacteria entering the medicine, the entire contents of the nebulizer should be used on the first time. **STORE THIS MEDICINE IN THE REFRIGERATOR,** between 36 and 46 degrees F (2 and 8 degrees C). Do not freeze. **UNOPENED POUCHES MAY BE STORED** at room temperature between 68 and 77 degrees F (20 and 25 degrees C) for up to 6 weeks. Store away from heat, moisture, and light. If this medicine is stored from room temperature for up to 6 weeks, **DO NOT USE THIS MEDICINE** if it is past the expiration date on the container. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **CONTINUE TO USE THIS MEDICINE** even if you feel well. Do not miss any doses. If you miss a dose of this medicine, skip the missed dose and go back to your regular dosing schedule. Do not use 2 doses at once.

CAUTIONS: **DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine or to formoterol. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescription or over-the-counter, check with your doctor or pharmacist. **LAB TESTS,** including lung function and blood potassium levels, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to keep all doctor and lab appointments. **THIS MEDICINE MAY CAUSE DIZZINESS.** This effect may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. Do not drive or perform other, possibly unsafe tasks until you know how you react to it. **THIS MEDICINE WILL NOT STOP SUDDEN SYMPTOMS OF COPD ONCE THEY HAVE ALREADY STARTED.** Be sure to carry a short-acting bronchodilator inhaler (eg, albuterol) with you at all times to treat any breathing problems that may occur between doses of this medicine (eg, severe or sudden onset of wheezing or shortness of breath). If you have any questions about which medicines stop sudden symptoms, check with your doctor or pharmacist. **IF YOU HAVE BEEN REGULARLY USING A SHORT-ACTING BRONCHODILATOR INHALER,** you may notice that you need to use about how to use it with this medicine. Short-acting bronchodilators are normally used with this medicine to treat breathing problems that may occur between doses. **THE RISK OF SERIOUS HEART PROBLEMS** (eg, irregular heartbeat) may be greater if you use this medicine in high doses. Do NOT use more than recommended dose or use more often than prescribed. **TELL YOUR DOCTOR AT ONCE** if you notice that your short-acting bronchodilator inhaler does not work as well, if you need to use it often, or if your breathing problems get worse. **CONTACT YOUR DOCTOR OR SEEK MEDICAL CARE RIGHT AWAY** if you have breathing problems that worsen quickly, or if you use your short-acting bronchodilator and do not get relief. **TALK WITH YOUR DOCTOR OR PHARMACIST** about all of your breathing medicines and how to use them. Do not start, stop, or change the dose of any

breathing medicine unless your doctor tells you to. **THE MEDICINE MAY SOMETIMES CAUSE SEVERE BREATHING PROBLEMS RIGHT AFTER YOU USE A DOSE.** If this happens, use your short-acting bronchodilator. **CONTACT YOUR DOCTOR OR SEEK OTHER MEDICAL CARE AT ONCE.** **THIS MEDICINE MAY RAISE YOUR BLOOD SUGAR.** High blood sugar may make you feel confused, drowsy, or thirsty. It can also make you flush, breathe faster, or have a fruit-like breath odor. If these symptoms occur, tell your doctor right away. **PREGNANCY and BREAST-FEEDING:** If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. It is not known if this medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while taking this medicine include: back pain; diarrhea; dry mouth; headache; nausea; nervousness; stuffy nose; tiredness; tremor; trouble sleeping; vomiting. If they continue or are bothersome, check with your doctor. **SEEK MEDICAL ATTENTION RIGHT AWAY** if any of these SEVERE side effects occur: chest pain; fast or irregular heartbeat; fever, chills, or persistent sore throat; leg swelling; new or worsening breathing problems (eg, increased chest tightness, coughing, shortness of breath, wheezing); severe or persistent headache, dizziness, tremor, or nervousness; severe or persistent muscle pain or cramps; symptoms of high blood sugar (eg, increased thirst, urination, or hunger; unusual weakness or drowsiness; confusion); trouble speaking. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, tongue, or throat; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include chest pain; fast or irregular heartbeat; severe or persistent dizziness; dry mouth; fatigue; headache; muscle pain or cramps; nausea; nervousness; trouble sleeping, or tremor; severe or persistent symptoms of high blood sugar (eg, increased thirst, urination, or hunger; drowsiness; flushing of the skin; confusion; fruit-like breath odor).

ADDITIONAL INFORMATION: If your symptoms do not improve or if they become worse, check with your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **USING THIS MEDICINE FOR AN EXTENDED PERIOD** of time may obtain refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 959810

KEEP OUT OF REACH OF CHILDREN. STORE IN SAFETY CONTAINER OR SECURE AREA.

<p>NELVA BRUNSTING 12850 Pinecroft, Houston, TX 770797517 (713)464-4321</p> <p>RX # 1496587-03328 DATE: 01/13/11</p>	<p>NELVA BRUNSTING 12850 Memorial Drive Houston, TX 77024 12850 Pinecroft, Houston, TX 770797517 (713)464-4321</p> <p>RX # 1496587-03328 DATE: 01/13/11</p>
<p>BROVANA 15MCG/2ML INH SOL 30X2ML QTY: 120 2 REFILLS BEFORE 12/05/11 Refill NDC:63402-0911-30</p> <p>\$ 527.99</p> <p>A. JAIN, MD MFG:SEPRACOR JDC/SSH/SSH/IKSC</p> <p>Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024 PH: (713)722-7247</p> <p>Customer Receipt</p>	<p>BROVANA 15MCG/2ML INH SOL 30X2ML QTY: 120 2 REFILLS BEFORE 12/05/11 Refill NDC:63402-0911-30</p> <p>\$ 527.99</p> <p>A. JAIN, MD MFG:SEPRACOR JDC/SSH/SSH/IKSC</p> <p>Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024 PH: (713)722-7247</p> <p>Duplicate Receipt</p>

Pharmacy use only

THU 4:15PM	BROVANA 15MCG/2ML INH SOL 30X2ML	QTY 120	LIQUID
Refill	63402-0911-30		
	REFRIG		

JDC/SSH/SSH/IKSC Med Guide

Brnstrung004471

Don't take a chance with your lungs — get a PARI Nebulizer cup now.



Did you know a PARI nebulizer was used to prove safety and efficacy of your inhaled medication? PARI nebulizers are the clinical choice, make them your choice.

The Benefits of PARI Reusable Nebulizers

- Short treatment times
- Consistent, efficient delivery of medication to the lungs
- Reusable, designed to last 6 months
- Easy to clean: boilable and dishwasher safe
- Cost effective

Specialists in effective inhalation



PARI LCP Plus used in clinical trial.

Continue to effectively manage your COPD by using a PARI Nebulizer.

Chronic obstructive pulmonary disease (COPD) is with you all of the time, even when you don't have symptoms. That is why it's important to use the PARI Nebulizer system, to deliver your medication to you effectively.


If you're not using a PARI nebulizer cup, you may not be getting the most benefit from your aerosol medication.

PARI reusable nebulizers feature breath enhanced technology resulting in increased aerosol delivery with decreased medication waste.

For more information contact PARI at 1.800.FAST.NEB (321.8632) or your Walgreens pharmacist. Visit www.PARI.com.

Talk to your doctor or pharmacist today about the PARI Reusable Nebulizer cup.

This message was developed and paid for by PARI Respiratory Equipment, Inc.



8
BR



1496587 0902 3 0062799 0


THU 4:15PM
\$527.99
EXPRESS PAY

NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

01/13/11
Refill

REFRIGERATE

- MEDICARE: Patient Not Eligible- Use New Plan or charge Cash
- Store TT 1 of 1
- Prescription Savings Club could save you \$117.29! Ask if you qualify.



**Personal
Prescription
Information**

LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.

Take advantage of these convenient services:

- **Touch Tone Refills**
Save time by using our automated system for a refill. Just dial the number on your prescription label.
- **Auto Refills**
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.



Visit us online at Walgreens.com

Thank you for choosing Walgreens!

To stop receiving sponsored info at this pharmacy, call 888-336-5744 or visit remove-me.net Use code:2875677 013 0320825

Brunsting004472

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
 12850 Memorial Drive
 Houston, TX 77024
 (713)722-7247

PATIENT NELVA BRUNSTING
BIRTH DATE 10/08/26
MEDICATION BROVANA 15MCG/2ML INH SOL 30X2ML
QUANTITY 120
DIRECTIONS INHALE 1 VIAL VIA NEBULIZER
TWICE DAILY

DOCTOR A. JAIN, MD
DRUG DESCRIPTION
 LIQUID
PATIENT ALLERGIES

INGREDIENT NAME: ARFORMOTEROL (ar-for-MOE-ter-of)

COMMON USES: This medicine is a long-acting beta-agonist bronchodilator used for long-term treatment of chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema. It may also be used to treat other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: WARNING: LONG-ACTING BETA-AGONISTS SUCH AS THIS MEDICINE HAVE BEEN RARELY ASSOCIATED WITH AN INCREASED RISK OF ASTHMA-RELATED DEATH. Long-acting beta-agonists should not be used in asthma patients without another long-term asthma-control medicine (eg, inhaled corticosteroids). This medicine has not been approved to treat asthma. Safety and effectiveness of this medicine in patients with asthma has not been confirmed. **SOME MEDICINES MAY INTERACT** with this medicine. **DO NOT TAKE THIS MEDICINE** if you are using another medicine that has a long-acting beta-agonist (eg, albuterol) in it. **TELL YOUR HEALTH CARE PROVIDER** if you are taking any other medicines, especially any of the following: corticosteroids (eg, prednisone) or a tricyclic antidepressant (eg, amitriptyline) within the last 14 days; theophylline; catechol-O-methyltransferase (COMT) inhibitors (eg, entacapone), monoamine oxidase inhibitors (MAOIs) (eg, phenelzine), tricyclic antidepressants (eg, amitriptyline) or beta-blockers (eg, propranolol) or heart rate, overexertion, or severe trouble sleeping. **Tell your doctor** if you have taken a MAOI (eg, phenelzine) or a tricyclic antidepressant (eg, amitriptyline) within the last 14 days. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you are having severe breathing problems (eg, sudden, severe onset or worsening of COPD symptoms such as chest tightness, cough, shortness of breath, wheezing), you have asthma and you are not currently using a long-term asthma-control medicine (eg, inhaled corticosteroids), or if you have asthma that is already well controlled with the use of a long-term asthma-control medicine. **THIS MEDICINE SHOULD NOT BE USED IN CHILDREN;** safety and effectiveness in children have not been confirmed. **Contact your doctor or pharmacist** if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine comes with a MEDICATION GUIDE approved by the U.S. Food and Drug Administration. Read it carefully each time you refill this medicine. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. **THIS MEDICINE SHOULD ONLY BE INHALED USING A NEBULIZER.** Do NOT inject or swallow it. A health care provider will teach you how to use the nebulizer. Be sure you know what type of nebulizer to use with this medicine, and how to use it. Follow the procedures you are taught when you use a dose. **Contact your health care provider** if you have any questions. **STORE THIS MEDICINE** in the original foil pouch in a dry place. Do not remove from the foil pouch or the vial until right before use. **IF THE MEDICINE CONTAINS PARTICLES** is cloudy or discolored, or if the vial is cracked or damaged in any way, do not use it. **DO NOT MIX THIS MEDICINE WITH OTHER MEDICINES** in your nebulizer machine. **TO USE THE MEDICINE:** twist open the top of the NEBULIZER RESERVOIR to the mouthpiece or face mask. **CONNECT THE MOUTHPIECE** in your mouth (or put on the face mask) and turn on the nebulizer. **INHALE** slowly, deeply, and evenly as possible until no more mist is formed in the nebulizer chamber (about 8 to 10 minutes). **CLEAN THE NEBULIZER** according to the instructions. Failure to properly clean the nebulizer could lead to bacteria entering the medicine. This may lead to an infection. To avoid bacteria entering the medicine, use the entire contents right after opening the vial for the first time. **STORE THE MEDICINE** in the refrigerator, between 36 and 46 degrees F (2 and 8 degrees C). Do not freeze. **UNOPENED POUCHES MAY BE STORED** at room temperature between 59 and 77 degrees F (15 and 25 degrees C) for up to 6 weeks. Store away from heat, moisture, and light. If the medicine is stored at room temperature, throw it away after 6 weeks. **DO NOT USE** this medicine if it is past the expiration date on the container. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **CONTINUE TO USE THIS MEDICINE** even if you feel well. Do not miss any doses. If you miss a dose of this medicine, skip the missed dose and go back to your regular dosing schedule. Do not use 2 doses at once.

CAUTIONS: DO NOT USE THIS MEDICINE if you are allergic to any ingredient in this medicine or to formoterol. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescription or over-the-counter, check with your doctor or pharmacist. **LAB TESTS,** including lung function and blood potassium levels, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to keep all doctor and lab appointments. **CAUTION: DIZZINESS CAN OCCUR** when this effect may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. **Do not drive or perform other possibly unsafe tasks until you know how you react to it.** **THIS MEDICINE WILL NOT STOP YOUR ASTHMA IF YOU HAVE ALREADY STARTED.** Be sure to carry a short-acting bronchodilator inhaler (eg, albuterol) with you at all times to treat any breathing problems that may occur between doses of this medicine (eg, severe or sudden onset of wheezing or shortness of breath). If you have any questions about which medicines you should use, check with your doctor or pharmacist. **Short-acting bronchodilators are normally used with this medicine** to treat breathing problems that may occur between doses. **THE RISK OF SERIOUS HEART PROBLEMS (eg, rapid heartbeat) may be greater** if you use this medicine in high doses. Do NOT use more than recommended dose or use more often than prescribed. **TELL YOUR DOCTOR AT ONCE** if you notice that your short-acting bronchodilator inhaler does not work as well, if you need to use it often, or if your breathing problems get worse. **CONTACT YOUR DOCTOR OR SEEK MEDICAL CARE RIGHT AWAY** if you have breathing problems that worsen quickly, or if you use your short-acting bronchodilator and do not get relief. **TALK WITH YOUR DOCTOR OR PHARMACIST** about all of your breathing medicines and how to use them. Do not start, stop, or change the dose of any

breathing medicine unless your doctor tells you to. **THE MEDICINE MAY SOMETIMES CAUSE SEVERE BREATHING PROBLEMS RIGHT AFTER YOU USE A DOSE.** If this happens, use your short-acting bronchodilator. **Contact your doctor or seek other medical care at once.** **THIS MEDICINE MAY RAISE YOUR BLOOD SUGAR.** High blood sugar may make you feel confused, drowsy, or thirsty. It can also make you flush, breathe faster, or have a fruit-like breath odor. If these symptoms occur, tell your doctor right away. **PREGNANCY AND BREAST-FEEDING:** If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. It is not known if this medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while taking this medicine include: back pain; diarrhea; dry mouth; headache; nausea; nervousness; stuffy nose; tiredness; tremor; trouble sleeping; vomiting. If they continue or are bothersome, check with your doctor. **SEEK MEDICAL ATTENTION RIGHT AWAY** if any of these SEVERE side effects occur: chest pain; fast or irregular heartbeat; fever; chills; or persistent sore throat; leg swelling; new or worsening breathing problems (eg, increased chest tightness, coughing, shortness of breath, wheezing); severe or persistent headache, dizziness, tremor, or nervousness; severe or persistent muscle pain or cramps; symptoms of high blood sugar (eg, increased thirst, urination, or hunger; unusual weakness or drowsiness; confusion); trouble speaking. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, tongue, or throat; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include chest pain; fast or irregular heartbeat; severe or persistent dizziness; dry mouth; fatigue; headache; muscle pain or cramps; nausea; nervousness; trouble sleeping; or tremors; severe or persistent symptoms of high blood sugar (eg, increased thirst, urination, or hunger; drowsiness; flushing of the skin; confusion; fruit-like breath odor).

ADDITIONAL INFORMATION: If your symptoms do not improve or if they become worse, check with your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** obtain refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain. WIC# 957918

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

NELVA BRUNSTING
 13630 Pinerock, Houston, TX 770797517
 (713)464-4391
RX # 1496587-03328 DATE: 02/18/11

BROVANA 15MCG/2ML INH SOL 30X2ML
 QTY: 120 1 REFILL BEFORE 12/05/11
 Refill NDC: 63402-0911-30 \$ 527.99

A. JAIN, MD
 MFG: SEPRACOR
 HMC / / / KDM

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247 Customer Receipt

NELVA BRUNSTING
 13630 Pinerock, Houston, TX 770797517
 (713)464-4391
RX # 1496587-03328 DATE: 02/18/11

BROVANA 15MCG/2ML INH SOL 30X2ML
 QTY: 120 1 REFILL BEFORE 12/05/11
 Refill NDC: 63402-0911-30 \$ 527.99

A. JAIN, MD
 MFG: SEPRACOR
 HMC / / / KDM

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247 Duplicate Receipt

Pharmacy use only

BROVANA 15MCG/2ML INH SOL 30X2ML QTY 120 LIQUID

TUE 12:00PM 63402-0911-30 Refill REFRIIG HMC / / / KDM Med Guide

Brnsting004467


Don't take a chance with your lungs — get a PARI Nebulizer cup now.



Did you know a PARI nebulizer was used to prove safety and efficacy of your inhaled medication? PARI nebulizers are the clinical choice, make them your choice.


The Benefits of PARI Reusable Nebulizers

- Short treatment times
- Consistent, efficient delivery of medication to the lungs
- Reusable, designed to last 6 months
- Easy to clean: boilable and dishwasher safe
- Cost effective



PARI LC® Plus used in clinical trial.

3544



Continue to effectively manage your COPD by using a PARI Nebulizer.

Chronic obstructive pulmonary disease (COPD) is with you all of the time, even when you don't have symptoms. That is why it's important to use the PARI Nebulizer system, to deliver your medication to you effectively.

If you're not using a PARI nebulizer cup, you may not be getting the most benefit from your aerosol medication.

PARI reusable nebulizers feature breath enhanced technology resulting in increased aerosol delivery with decreased medication waste.



Talk to your doctor or pharmacist today about the PARI Reusable Nebulizer cup.

This message was developed and paid for by PARI Respiratory Equipment, Inc.

OCT-2875 677 049 0537281

INFO: 0806.63402091130

3544

8 BR



1486587 2203 3 0062798 0

NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

- MEDICARE: Patient Not Eligible- Use New Plan or charge Cash
 - Store TT 1 of 1
 - Prescription Savings Club could save you \$117.29!
- Ask if you qualify.

TUE 12:00PM
\$527.99
EXPRESS PAY

02/18/11

Refill

REFRIGERATE
MED GUIDE



Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.

Take advantage of these convenient services:

- **Auto Refills**
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.
- **Touch Tone Refills**
Save time by using our automated system for a refill. Just dial the number on your prescription label.

Walgreens

The Pharmacy America Trusts • Since 1901™

Visit us online at Walgreens.com


Thank you for choosing Walgreens!

To stop receiving sponsored info at this pharmacy, call 888-336-5744 or visit remove-me.net Use code:2875677 049 0537281

Brunsting004468

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
 12850 Memorial Drive
 Houston, TX 77024
 (713)722-7247

PATIENT NELVA BRUNSTING	DOCTOR A. JAIN, MD	DRUG DESCRIPTION
BIRTH DATE 10/08/26		
MEDICATION AZITHROMYCIN 250MG TABLETS	PATIENT ALLERGIES	
QUANTITY 30		
DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY DAY		

PINK
 FRONT: 93
 BACK: 7146

INGREDIENT NAME: AZITHROMYCIN
 (ay-ZITH-roe-MYE-sin)

COMMON USES: This medicine is a macrolide antibiotic used to treat bacterial infections.

BEFORE USING THIS MEDICINE: Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking dofetilide, nilotinib, propafenone, pimozide, or tetrabenazine. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking antiarrhythmics (eg, disopyramide), anticoagulants (eg, warfarin), arsenic, astemizole, carbamazepine, cisapride, digoxin, domperidone, maprotiline, methadone, nelfinavir, cyclosporine, ergot alkaloids (eg, ergotamine), paliperidone, phenytoin, quinolones (eg, levofloxacin), rifampin, terfenadine, theophylline, triazolam, tyrosine kinase inhibitors (eg, dasatinib), or medicines that may affect your heartbeat. Ask your doctor if you are unsure if any of the medicines you are taking may affect your heartbeat. Inform your doctor of any other medical conditions including irregular heartbeat, kidney problems, liver problems, myasthenia gravis, allergies, pregnancy or breastfeeding. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine may be taken on an empty stomach or with food. DO NOT TAKE THIS MEDICINE within 1 hour before or 2 hours after aluminum- or magnesium-containing antacids. STORE THIS MEDICINE at room temperature, away from heat and light. TO CLEAR UP YOUR INFECTION COMPLETELY, continue taking this medicine for the full course of treatment even if you feel better in a few days. Do not miss any doses. Taking this medicine at the same time each day will make it easier to remember. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. If you miss a dose, do not take 2 doses at once.

CAUTIONS: DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it, to other macrolide antibiotics (such as erythromycin), to ketolide antibiotics (such as telithromycin), or if you are allergic to any ingredient in this product. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to erythromycin or any macrolide or ketolide antibiotic. A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine, contact your doctor or pharmacist. IF YOU EXPERIENCE difficulty breathing; tightness of chest; swelling of eyelids, face, or lips; or if you develop a rash or hives, tell your doctor immediately. Do not take any more of this medicine unless your doctor tells you to do so. This medicine may cause drowsiness, dizziness, or lightheadedness. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. THIS

MEDICINE MAY CAUSE increased sensitivity to the sun. Avoid exposure to the sun, sunlamps, or tanning booths until you know how you react to this medicine. Use a sunscreen or protective clothing if you must be outside for a prolonged period. MILD DIARRHEA IS COMMON with antibiotic use. However, a more serious form of diarrhea (pseudomembranous colitis) may rarely occur. This may develop while you use the antibiotic or within several months after you stop using it. Contact your doctor right away if stomach pain or cramps, severe diarrhea, or bloody stools occur. Do not treat diarrhea without first checking with your doctor. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. FOR WOMEN: IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. IT IS UNKNOWN IF THIS MEDICINE IS EXCRETED in breast milk. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS, that may go away during treatment, include mild diarrhea, headache, nausea, or stomach pain or upset. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience vomiting, changes in hearing, or ringing in the ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience chest pain; swelling of your hands, legs, face, lips, eyes, throat, or tongue; difficulty swallowing, speaking, or breathing; eye or vision problems; fainting; hearing loss; irregular heartbeat; muscle weakness; reddened, blistered, or swollen skin; seizures; severe stomach pain, cramps, or diarrhea; bloody stools; unusual vaginal itching, odor, or discharge; or yellowing of the eyes or skin. An allergic reaction to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, dizziness, trouble breathing, or unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include nausea, vomiting, stomach upset, and diarrhea.

ADDITIONAL INFORMATION: DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957818

NELVA BRUNSTING
 13630 Pinerock, Houston, TX 770797517
 (713)464-4391
RX # 1494789-03328 DATE: 03/02/11

AZITHROMYCIN 250MG TABLETS
 QTY: 30 NO REFILLS - DR. AUTH REQUIRED
 Refill NDC: 00093-7146-56
 Retail Price: \$195.79 Your Insurance Saved You: \$190.79 \$ 5.00

A. JAIN, MD PLAN: PAIDMPD
 MFG: TEVA GROUP# CMD3896
 XXX / / KMN/KMN CLAIM REF# FKRWQHL

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247 Customer Receipt

NELVA BRUNSTING
 13630 Pinerock, Houston, TX 770797517
 (713)464-4391
RX # 1494789-03328 DATE: 03/02/11

AZITHROMYCIN 250MG TABLETS
 QTY: 30 NO REFILLS - DR. AUTH REQUIRED
 Refill NDC: 00093-7146-56
 Retail Price: \$195.79 Your Insurance Saved You: \$190.79 \$ 5.00

A. JAIN, MD PLAN: PAIDMPD
 MFG: TEVA GROUP# CMD3896
 XXX / / KMN/KMN CLAIM REF# FKRWQHL

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247 Duplicate Receipt

Pharmacy use only

WED 1:30PM AZITHROMYCIN 250MG TABLETS
 Refill 00093-7146-56
 ALPHA

QTY 30
 10 DRAM



PINK
 FRONT: 93
 BACK: 7146

XXX / / KMN/KMN

Brunsting004465

AABrunsting.Financials003958

Ask if an antibiotic is causing her diaper rash.

Without leaving her side.



New Pharmacy Chat at Walgreens.com

helps you find immediate answers anytime, anywhere.

Chat live with an expert from our pharmacy team about:

- Your personal prescription questions
- Over-the-counter product and prescription interactions
- Treatments for common ailments



Go to Walgreens.com/pharmacychat today.

It's confidential and secure!

Walgreens

Arm yourself[®] for the ones you love



No appointment necessary.

Flu shots* every day.

Register for your flu shot today at Walgreens.com/flu.

We bill many insurance plans directly, including Medicare.

*Prices may vary for certain forms of the flu vaccine. Vaccines subject to availability. State, age and health condition-related restrictions may apply. See pharmacy for details.

Walgreens

8 BR




NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

- Store TT 3 of 6
- Your Insurance Saved You: \$190.79

WED 1:30PM
\$5.00
EXPRESS PAY

03/02/11
Refill



Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- **Auto Refills**
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

Walgreens
The Pharmacy America Trusts • Since 1901[®]

Visit us online at Walgreens.com

Thank you for choosing Walgreens!

Brunsting004466

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
12850 Memorial Drive
Houston, TX 77024
(713)722-7247

PATIENT NELVA BRUNSTING
BIRTH DATE 10/08/26
MEDICATION SERTRALINE 50MG TABLETS
QUANTITY 30
DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY DAY

DOCTOR A. JAIN, MD

DRUG DESCRIPTION



BLUE
FRONT: G 4900
BACK: 50MG

INGREDIENT NAME: SERTRALINE (SER-tra-leen)

COMMON USES: This medicine is a selective serotonin reuptake inhibitor (SSRI) used to treat depression, panic disorder, obsessive-compulsive disorder (OCD), post-traumatic stress disorder (PTSD), social anxiety disorder (social phobia), and a severe form of premenstrual syndrome called premenstrual dysphoric disorder (PMDD). It may also be used for other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: WARNING: Antidepressants may increase the risk of suicidal thoughts or actions in children, teenagers, and young adults. However, depression and certain other mental problems may also increase the risk of suicide. Talk with the patient's doctor to be sure that the benefits of using this medicine outweigh the risk and caregivers must closely watch patients who take this medicine. It is important to keep in close contact with the patient's doctor. Tell the doctor right away if the patient has symptoms like worsened depression, suicidal thoughts, or changes in behavior. Discuss any questions with the patient's doctor. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking a fenfluramine derivative (eg, dexfenfluramine); nefazodone; pimozide; sibutramine; or thioridazine. **DO NOT TAKE THIS MEDICINE** if you are taking or have taken a monoamine oxidase inhibitor (MAOI) (eg, phenelzine), selegiline, or St. John's wort within the last 14 days. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking anorexiants (eg, phentermine); linezolid; metoclopramide; serotonin 5-HT1 receptor agonists (eg, sumatriptan); trazodone; anticoagulants (eg, warfarin); aspirin; nonsteroidal anti-inflammatory drugs (NSAIDs) (eg, ibuprofen); diuretics (eg, furosemide, hydrochlorothiazide); tramadol; phenothiazines (eg, chlorpromazine); carbamazepine; cyproheptadine; antipyrazole; clozapine; digoxin; flecainide; lithium; phenytoin; propafenone; risperidone; tricyclic antidepressants (eg, amitriptyline); or valproic acid (eg, valproate). **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including if you have a history of seizures; heart problems; liver problems; stomach or bowel bleeding; metabolism problems; allergies; pregnancy; or breast-feeding. Tell your doctor if you or a family member has a history of bipolar disorder (manic-depression), other mental or mood problems, suicidal thoughts or attempts, or alcohol or substance abuse. Tell your doctor if you are dehydrated, have low blood sodium levels, drink alcohol, or if you will be having electroconvulsive therapy (ECT). Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for taking this medicine provided by your doctor. This medicine has a **MEDICATION GUIDE** approved by the U.S. Food and Drug Administration. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. **TAKE THIS MEDICINE** by mouth with or without food. **STORE THIS MEDICINE** at 77 degrees F (25 degrees C) away from heat, moisture, and light. Brief storage at temperatures between 59 and 86 degrees F (15 and 30 degrees C) is permitted. Do not store in the bathroom. **CONTINUE TO TAKE THIS MEDICINE** even if you feel well. Do not miss any doses. Taking this medicine at the same time each day will help you remember to take it. **DO NOT SUDDENLY STOP TAKING THIS MEDICINE** without checking with your doctor. Side effects may occur. They may include mental or mood change, numbness or tingling of the skin, dizziness, confusion, headache, trouble sleeping, or unusual tiredness. You will be closely monitored when you start this medicine and whenever a change in dose is made. **IF YOU MISS A DOSE OF THIS MEDICINE**, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **DO NOT** take 2 doses at once.

CAUTIONS: **DO NOT USE THIS MEDICINE** if you are allergic to any ingredient in this medicine. **THIS MEDICINE MAY CAUSE DROWSINESS OR DIZZINESS.** These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. **DO NOT DRIVE OR PERFORM OTHER POSSIBLY UNSAFE TASKS** until you know how you react to it. **DO NOT DRINK ALCOHOL** while you are taking this medicine. Check with your doctor before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers) while you are taking this medicine; it may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. Several weeks may pass before your symptoms improve. **DO NOT TAKE MORE THAN THE RECOMMENDED DOSE**, change your dose, or use this medicine for longer than prescribed without checking with your doctor. **IF YOUR DOCTOR**

TELLS YOU TO STOP TAKING THIS MEDICINE, you will need to wait for several weeks before beginning to take certain other medicines (eg, MAOIs, nefazodone). Ask your doctor when you should start to take your new medicines after you have stopped taking this medicine. **SEROTONIN SYNDROME** and **NEUROLEPTIC MALIGNANT SYNDROME (NMS)** are possibly fatal syndromes that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans", MAOIs, antipsychotics). Symptoms of these syndromes may include blood pressure changes; agitation; confusion; hallucinations; other mental or mood changes; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; rigid muscles; and nausea, vomiting, or diarrhea. **CONTACT YOUR DOCTOR** at once if you have any of these symptoms. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES**, either prescription or over-the-counter, check with your doctor or pharmacist. Caution is advised when using this medicine in the **ELDERLY**; they may be more sensitive to its effects, especially low blood sodium levels. Caution is advised when using this medicine in **CHILDREN**; they may be more sensitive to its effects, especially increased risk of suicidal thoughts or actions. **THIS MEDICINE MAY CAUSE WEIGHT CHANGES; CHILDREN AND TEENAGERS** may need regular weight and growth checks while they take this medicine. **FOR MEN: THIS MEDICINE MAY RARELY CAUSE A** painful erection. This could happen even when you are not having sex. If this is not treated right away, it could lead to permanent sexual problems such as impotence. Contact your doctor right away if this happens. **FOR WOMEN: THIS MEDICINE MAY CAUSE HARM TO THE FETUS** if it is used during the last 3 months of pregnancy. **IF YOU BECOME PREGNANT**, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. **IT IS NOT KNOWN IF THIS MEDICINE IS FOUND IN breast milk.** **IF YOU ARE OR WILL BE BREAST-FEEDING** while you use this medicine, check with your doctor. Discuss any possible risks to your baby.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while taking this medicine include anxiety; constipation; decreased sexual desire or ability; diarrhea; dizziness; drowsiness; dry mouth; increased sweating; loss of appetite; nausea; nervousness; stomach upset; tiredness; trouble sleeping; vomiting; or weight loss. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience bizarre behavior; black or bloody stools; chest pain; decreased bladder control; exaggerated reflexes; fast or irregular heartbeat; fever; hallucinations; loss of coordination; new or worsening agitation; panic attacks; aggressiveness, impulsiveness, irritability, hostility, exaggerated feeling of well-being, restlessness, or inability to sit still; persistent or severe ringing in the ears; persistent, painful erection; red, swollen, blistered, or peeling skin; seizures; severe or persistent anxiety or trouble sleeping; stomach pain; suicidal thoughts or attempts; tremor; unusual bruising or bleeding; unusual or severe mental or mood changes; vision changes; or worsening of depression. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include coma; fainting; fast, slow, or irregular heartbeat; hair loss; hallucinations; seizures; severe or persistent dizziness; drowsiness, diarrhea, nausea, or vomiting; or tremor.

ADDITIONAL INFORMATION: **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME**, obtain refills before your supply runs out.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

NELVA BRUNSTING

13630 Pinrock, Houston, TX 770797517
(713)464-4391

RX # 1515376-03328

DATE: 03/02/11

SERTRALINE 50MG TABLETS

QTY: 30 1 REFILL BEFORE 02/02/12

Refill NDC: 59762-4900-05

Retail Price: \$29.99 Your Insurance Saved You: \$24.99

\$ 5.00

A. JAIN, MD
MFG: GREENSTONE
XXX/KMN/KMN / KMN

PLAN: PAIDMPD
GROUP# CMD3896
CLAIM REF# OXHXMT3

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
PH: (713)722-7247

Customer Receipt

NELVA BRUNSTING

13630 Pinrock, Houston, TX 770797517
(713)464-4391

RX # 1515376-03328

DATE: 03/02/11

SERTRALINE 50MG TABLETS

QTY: 30 1 REFILL BEFORE 02/02/12

Refill NDC: 59762-4900-05

Retail Price: \$29.99 Your Insurance Saved You: \$24.99

\$ 5.00

A. JAIN, MD
MFG: GREENSTONE
XXX/KMN/KMN / KMN

PLAN: PAIDMPD
GROUP# CMD3896
CLAIM REF# OXHXMT3

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

SERTRALINE 50MG TABLETS

59762-4900-05

CELL 29

QTY 30

20 DRAM



BLUE
FRONT: G 4900
BACK: 50MG

XXX/KMN/KMN / KMN

Med Guide

Brunsting004463

AABrunsting.Financials003960

Ask if grapefruit juice affects your medication.

From the breakfast table.



New Pharmacy Chat at Walgreens.com

helps you find immediate answers anytime, anywhere.

Chat live with an expert from our pharmacy team about:

- Your personal prescription questions
- Over-the-counter product and prescription interactions
- Treatments for common ailments




Go to Walgreens.com/pharmacychat today.

It's confidential and secure!

Walgreens

3883

Arm yourself® for the ones you love



No appointment necessary.

Flu shots* every day.

Register for your flu shot today at Walgreens.com/flu.

We bill many insurance plans directly, including Medicare.

*Prices may vary for certain forms of the flu vaccine. Vaccines subject to availability. State, age and health condition-related restrictions may apply. See pharmacy for details.

Walgreens

OTC: 2875 67 061 0615664

INFC: 0908 99762490005

5883

8 BR




NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

- Store TT 5 of 6
- Your Insurance Saved You: \$24.99

WED 1:30PM
\$5.00
EXPRESS PAY

03/02/11
Refill
MED GUIDE



Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- Auto Refills
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

Walgreens

The Pharmacy America Trusts • Since 1901™

Visit us online at Walgreens.com

Thank you for choosing Walgreens!

Brunsting004464

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
 12850 Memorial Drive
 Houston, TX 77024
 (713)722-7247

PATIENT	NELVA BRUNSTING	DOCTOR	A. JAIN, MD	DRUG DESCRIPTION	LIQUID
BIRTH DATE	10/08/26				
MEDICATION	BROVANA 15MCG/2ML INH SOL 30X2ML	PATIENT ALLERGIES			
QUANTITY	120				
DIRECTIONS	INHALE 1 VIAL VIA NEBULIZER TWICE DAILY				

INGREDIENT NAME: ARFORMOTEROL (ar-for-MOE-ter-ole)

COMMON USES: This medicine is a long-acting beta-agonist bronchodilator used for long-term treatment of chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema. It may also be used to treat other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: WARNING: LONG-ACTING BETA-AGONISTS SUCH AS THIS MEDICINE HAVE BEEN RARELY ASSOCIATED WITH AN INCREASED RISK OF ASTHMA-RELATED DEATH. Long-acting beta-agonists should not be used in asthma patients without another long-term asthma-control medicine (eg, inhaled corticosteroids). This medicine has not been approved to treat asthma. Safety and effectiveness of this medicine in patients with asthma have not been confirmed. **SOME MEDICINES MAY INTERACT** with this medicine. **DO NOT TAKE THIS MEDICINE** if you are using another medicine that has a long-acting beta-agonist (eg, salmeterol) in it. **TELL YOUR HEALTH CARE PROVIDER** if you are taking any other medicines, especially any of the following: corticosteroids (eg, prednisone), diuretics (eg, furosemide, hydrochlorothiazide), xanthines (eg, theophylline), carbonic dehydratase (CDH) inhibitors (eg, etacrynic acid), monoamine oxidase inhibitors (MAOIs) (eg, phenelzine), tricyclic antidepressants (eg, amitriptyline), or any other medicine. **ASK YOUR HEALTH CARE PROVIDER** if this medicine may interact with other medicines you take. **DO NOT START OR STOP ANY MEDICINE** without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including a history of other breathing problems (eg, asthma), diabetes, heart problems (eg, fast or irregular heartbeat, heart blood vessel problems), liver problems, high blood pressure, low blood potassium levels, seizures, an overactive thyroid, or you have high blood or urine ketone levels. **ALLERGY, PREGNANCY, OR BREAST-FEEDING:** Tell your doctor if you have recently been to an emergency room for breathing problems; have a history of frequent hospitalizations for breathing problems; or have had an unusual reaction to a sympathomimetic medicine (eg, albuterol, pseudoephedrine), such as fast or irregular heartbeat, overexcitement, or severe trouble sleeping. **TELL YOUR DOCTOR** if you have taken a MAOI (eg, phenelzine) or a tricyclic antidepressant (eg, amitriptyline) within the last 14 days. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you are having severe breathing problems (eg, sudden, severe onset or worsening of COPD symptoms such as chest tightness, cough, shortness of breath, wheezing), you have asthma and you are not currently using asthma-control medicine (eg, inhaled corticosteroids), or if you have asthma that is already well controlled with the use of a long-term asthma-control medicine. **THIS MEDICINE SHOULD NOT BE USED IN CHILDREN;** safety and effectiveness in children have not been confirmed. **Contact your doctor or pharmacist** if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine comes with a MEDICATION GUIDE approved by the U.S. Food and Drug Administration. Read it carefully each time you take this medicine. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. **THIS MEDICINE SHOULD ONLY BE INHALED USING A NEBULIZER.** Do not inject or swallow it. A health care provider will teach you how to use the nebulizer. Be sure you know what type of nebulizer to use with this medicine and how to use it. Follow the procedures you are taught when you use a dose. **Contact your health care provider** if you have any questions. **STORE THIS MEDICINE** in the original foil pouch in a dry place. **DISCONNECT THE NEBULIZER** from the foil pouch of the vial unit right before use. **IF THE MEDICINE CONTAINS PARTICLES,** is cloudy or discolored, or if the vial is cracked or damaged in any way, do not use it. **DO NOT MIX THIS MEDICINE WITH OTHER MEDICINES** in your nebulizer machine. **TO USE THIS MEDICINE,** twist open the top of the vial and the entire contents into the nebulizer reservoir. **CONNECT THE NEBULIZER RESERVOIR** to the mouthpiece or face mask. **CONNECT THE NEBULIZER** to the compressor. Sit in a comfortable, upright position. **PLACE THE MOUTHPIECE OR FACE MASK** in your mouth and turn on the nebulizer. **BREATHE AS CALMLY,** deeply, and evenly as possible until no more mist is formed in the nebulizer (about 5 to 10 minutes). **CLEAN THE NEBULIZER** according to the instructions. Failure to properly clean the nebulizer could lead to bacteria entering the medicine. This may lead to an infection. To avoid bacteria entering the medicine, use the spray canister immediately after opening the vial for the first time. **STORE THIS MEDICINE** in the REFRIGERATOR, between 36 and 46 degrees F (2 and 8 degrees C). Do not freeze. **UNOPENED POUCHES** may be stored at room temperature between 68 and 77 degrees F (20 and 25 degrees C) for up to 6 weeks. Store away from heat, moisture, and light. If this medicine is stored at room temperature, throw it away after 6 weeks. **DO NOT USE** this medicine if it is past the expiration date on the container. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **CONTINUE TO USE THIS MEDICINE** even if you feel well. Do not miss any doses. If you miss a dose of this medicine, skip the missed dose and go back to your regular dosing schedule. Do not use 2 doses at once.

CAUTIONS: DO NOT USE THIS MEDICINE if you are allergic to any ingredient in this medicine or to formoterol. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either by prescription or over-the-counter, check with your doctor or pharmacist. **LAB TESTS,** including lung function and blood potassium levels, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to keep all doctor or lab appointments. **THIS MEDICINE MAY CAUSE DIZZINESS.** This effect may be worse if you take it with alcohol or other medicines. Use this medicine with caution. **Do not drive or perform other possibly unsafe tasks** until you know how you react to it. **THIS MEDICINE WILL NOT STOP SUDDEN SYMPTOMS OF COPD** ONCE THEY HAVE ALREADY STARTED. Be sure to carry a short-acting bronchodilator inhaler (eg, albuterol) with you at all times to treat any breathing problems that may occur between doses of this medicine (eg, severe or sudden onset of wheezing or shortness of breath). If you have any questions about which medicines stop sudden symptoms, check with your doctor or pharmacist. **IF YOU HAVE BEEN TOLD EARLY USING A SHORT-ACTING BRONCHODILATOR INHALER,** talk with your doctor about how to use it with this medicine. Short-acting bronchodilators are normally used with this medicine to treat breathing problems that may occur between doses. **THE RISK OF SERIOUS HEART PROBLEMS** (eg, irregular heartbeat) may be greater if you use this medicine in high doses. Do not use more than recommended dose or use more often than prescribed. **TELL YOUR DOCTOR AT ONCE** if you notice that your short-acting bronchodilator inhaler does not work as well as you need to use it often, or if your breathing problems get worse. **CONTACT YOUR DOCTOR OR SEEK MEDICAL CARE RIGHT AWAY** if you have breathing problems that worsen quickly. Do not stop using your short-acting bronchodilator and do not get relief. **TALK WITH YOUR DOCTOR OR PHARMACIST** about all of your breathing medicines and how to use them. Do not start, stop, or change the dose of any

breathing medicine unless your doctor tells you to. **THE MEDICINE MAY SOME TIMES CAUSE SEVERE BREATHING PROBLEMS RIGHT AFTER YOU USE A DOSE.** If this happens, use your short-acting bronchodilator. **Contact your doctor or seek other medical care at once.** **THIS MEDICINE MAY RAISE YOUR BLOOD SUGAR.** High blood sugar may make you feel confused, drowsy, or thirsty. It can also make you flush, breathe faster, or have a fruit-like breath odor. If these symptoms occur, tell your doctor right away. **PREGNANCY AND BREAST-FEEDING:** If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. It is not known if this medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby.

POSSIBLE SIDE EFFECTS; SIDE EFFECTS that may occur while taking this medicine include: back pain; diarrhea; dry mouth; headache; nausea; nervousness; stuffy nose; tiredness; tremor; trouble sleeping; vomiting. If they continue or are bothersome, check with your doctor. **SEEK MEDICAL ATTENTION RIGHT AWAY** if any of these SEVERE side effects occur: chest pain; fast or irregular heartbeat; fever, chills, or persistent sore throat; leg swelling; new or worsening breathing problems (eg, increased chest tightness, coughing, shortness of breath, wheezing); severe or persistent headache, dizziness, tremor, or nervousness; severe or persistent muscle pain or cramps; symptoms of high blood sugar (eg, increased thirst, urination, or hunger; unusual weakness or drowsiness; confusion); trouble speaking. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, tongue, or throat; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include chest pain; fast or irregular heartbeat; severe or persistent dizziness, dry mouth, fatigue, headache, muscle pain or cramps, nausea, nervousness, trouble sleeping, or tremors; severe or persistent symptoms of high blood sugar (eg, increased thirst, urination, or hunger; drowsiness; flushing of the skin; confusion; fruit-like breath odor).

ADDITIONAL INFORMATION: If your symptoms do not improve or if they become worse, check with your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **IF USING THIS MEDICINE** in an extended-release or timed-release container, check with your pharmacist about all of your breathing medicines and how to dispose of unused medicine.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush medications or pour down a sink or drain.

WICK 957918

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

NELVA BRUNSTING
 13630 Pinerock, Houston, TX 770797517
 (713)464-4391

RX # 1540088-03328 DATE: 04/21/11

BROVANA 15MCG/2ML INH SOL 30X2ML
 QTY: 120 2 REFILLS BEFORE 04/21/12
 Copy NDC: 63402-0911-30

\$ 527.99

A. JAIN, MD
 MFG:SEPRACOR
 XXX/RJW/KHT/ /PBP

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247

Customer Receipt

NELVA BRUNSTING
 13630 Pinerock, Houston, TX 770797517
 (713)464-4391

RX # 1540088-03328 DATE: 04/21/11

BROVANA 15MCG/2ML INH SOL 30X2ML
 QTY: 120 2 REFILLS BEFORE 04/21/12
 Copy NDC: 63402-0911-30

\$ 527.99

A. JAIN, MD
 MFG:SEPRACOR
 XXX/RJW/KHT/ /PBP

Walgreens

12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247

Duplicate Receipt

Pharmacy use only

THU 5:16PM
 Copy

BROVANA 15MCG/2ML INH SOL 30X2ML
 QTY 120
 LIQUID

63402-0911-30
 REFRIG

XXX/RJW/KHT/ /PBP

Med Guide

Brunsting004457

AABrunsting.Financials003962

Save up to 30% on your prescriptions

Enroll today for only \$19.95!

The AARP® Prescription Discount Program from Walgreens saves you up to 30 percent on all your FDA-approved prescriptions. This includes generic, brand name and specialty medications.



How Does It Work?

Your AARP Prescription Discount Program card can be used for drugs not covered by your insurance plan or if you have no prescription insurance. Simply present your AARP Prescription Discount Program card to your Walgreens pharmacist — after any other insurance card you may also have — when you fill or refill a prescription.

Note: This program is not a prescription drug insurance plan or a Medicare Part D plan, and does not replace such coverage.

How Do I Enroll?

You must be an AARP member to enroll. We can enroll you into both programs with one easy call. Just dial 1-877-4AARP19 (1-877-422-7719). You can also enroll online by visiting us at www.aarp-pharmacy.com.

What If I Have Questions?

Visit aarp-pharmacy.com or call the Walgreens Customer Care Center toll free, 24/7 at 1-877-4AARP19 (1-877-422-7719), or TTY 1-800-925-0178.

8
BR



1540088 0501 3 0052799 5

NELVA BRUNSTING


13630 Pinerock
Houston, TX 77079-7517
(713)464-4381

- PAIDMPP: Prior Authorization Required
- Prescription Savings Club could save you \$117.29!
- Ask if you qualify.

THU 5:16PM
\$527.99
EXPRESS PAY

04/21/11
Copy

REFRIGERATE
MED GUIDE



Personal Prescription Information

LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.

Take advantage of these convenient services:

- **Touch Tone Refills**
Save time by using our automated system for a refill. Just dial the number on your prescription label.
- **Auto Refills**
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

Walgreens
The Pharmacy America Trusts • Since 1901™

Visit us online at Walgreens.com

Thank you for choosing Walgreens!

OPT: 2875 677 116 0935995

INFO: 0916 63402091130


57616

AA Brunsting.Financials003963

Brunsting004458

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
 12850 Memorial Drive
 Houston, TX 77024
 (713)722-7247

PATIENT NELVA BRUNSTING	DOCTOR A. JAIN, MD	DRUG DESCRIPTION
BIRTH DATE 10/08/26		
MEDICATION ETHAMBUTOL 400MG TABLETS	PATIENT ALLERGIES	
QUANTITY 90		WHITE
DIRECTIONS TAKE 3 TABLETS BY MOUTH EVERY DAY		FRONT: L U
		BACK: C32

INGREDIENT NAME: ETHAMBUTOL (e-THAM-byoo-tole)

COMMON USES: This medicine is an antibacterial used to treat tuberculosis (TB).

BEFORE USING THIS MEDICINE: INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. Inform your doctor of any other medical conditions, allergies, pregnancy, or breast-feeding.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. STORE THIS MEDICINE at room temperature in a tightly-closed container, away from heat and light. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

CAUTIONS: DO NOT STOP USING THIS MEDICINE without first checking with your doctor. THIS MEDICINE MAY CAUSE dizziness. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. HAVE REGULAR EYE EXAMINATIONS while you are taking this medicine even if you do not notice changes in your vision. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS, that may go away during treatment, include nausea, headache, or dizziness. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience changes in vision, vomiting, skin rash, itching, fever, confusion, or numbness or tingling of extremities. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately.

ADDITIONAL INFORMATION: If your symptoms do not improve within 2 to 3 weeks, or if they become worse, check with your doctor. DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

NELVA BRUNSTING
 13630 Pinrock, Houston, TX 770797517
 (713)464-4391
RX # 1534700-03328 DATE: 04/29/11

ETHAMBUTOL 400MG TABLETS
 QTY: 90 1 REFILL BEFORE 04/05/12
 Refill NDC: 68180-0281-01
 Retail Price: \$153.59 Your Insurance Saved You: \$148.59 \$ 5.00

A. JAIN, MD PLAN: PAID/MPD
 MFG: LUPIN GROUP# CMD3896
 XXX/PBP/PBP/ /PBP CLAIM REF# 7C1HH79

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247 Customer Receipt

NELVA BRUNSTING
 13630 Pinrock, Houston, TX 770797517
 (713)464-4391
RX # 1534700-03328 DATE: 04/29/11

ETHAMBUTOL 400MG TABLETS
 QTY: 90 1 REFILL BEFORE 04/05/12
 Refill NDC: 68180-0281-01
 Retail Price: \$153.59 Your Insurance Saved You: \$148.59 \$ 5.00

A. JAIN, MD PLAN: PAID/MPD
 MFG: LUPIN GROUP# CMD3896
 XXX/PBP/PBP/ /PBP CLAIM REF# 7C1HH79


Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024
 PH: (713)722-7247 Duplicate Receipt

Pharmacy use only

SUN 11:00AM ETHAMBUTOL 400MG TABLETS QTY 90

Refill 68180-0281-01

ALPHA

 WHITE
 FRONT: L U
 BACK: C32

XXX/PBP/PBP/ /PBP

Brunsting004461



Take your antibiotics the right way.

- Precisely follow usage directions.
- Do not skip doses.
- Do not share them with others.
- Finish the prescription even if you feel better.
- Do not save them for future use.

Why is this checklist so important?

Using an antibiotic the wrong way can make infections stronger and harder to treat. You can prevent this problem by getting smart about antibiotics.

For more information talk to your Walgreens pharmacist. Or call 1-800-CDC-INFO or visit www.cdc.gov/getsmart.

5782

Do you have a higher risk of getting pneumonia?

Are you:

- Age 19 through 64 and smoke or have asthma?
- Age 64 or younger and have diabetes, heart disease, lung disease, leukemia, lymphoma, Hodgkin's disease, kidney problems, HIV or other condition that lowers the body's resistance to infection?
- Age 64 or younger and are taking a drug or treatment that lowers the body's resistance to infection, such as long-term steroids, certain cancer drugs or radiation therapy?
- Age 65 or older?

If you answered, "yes" to any of these questions, the Centers for Disease Control & Prevention (CDC) recommends that you receive a pneumonia vaccination.

**Talk to your Walgreens pharmacist
to get vaccinated today!**

No out-of-pocket cost for Medicare Part B beneficiaries*!

There's a way to stay well.

*Medicare Part B generally covers the pneumonia vaccine once per beneficiary. Vaccine subject to availability. State, age and health condition-related restrictions may apply. See pharmacy for details.

OFF: 2875 677 119 0964505

INFO: 0977 681 80028101

5782

8
BR



NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

- Store TT 1 of 5
- Your Insurance Saved You: \$148.59

SUN 11:00AM
\$5.00
EXPRESS PAY

04/29/11
Refill



**Personal
Prescription
Information**

LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- Auto Refills
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

The Pharmacy America Trusts • Since 1901*


Visit us online at Walgreens.com

Thank you for choosing Walgreens!

Brunsting004462

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
12850 Memorial Drive
Houston, TX 77024
(713)722-7247

PATIENT NELVA BRUNSTING
BIRTH DATE 10/08/26
MEDICATION METOPROLOL TARTRATE 50MG TABLETS
QUANTITY 60
DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY 12 HOURS
DOCTOR A. JAIN, MD
PATIENT ALLERGIES
DRUG DESCRIPTION

PINK
FRONT: m 32

INGREDIENT NAME: METOPROLOL (me-TOE-proe-IOLE)

COMMON USES: This medicine is a beta-adrenergic blocking agent (beta-blocker) used to treat high blood pressure. It may be used alone or with other medicines. It is also used for the long-term treatment of chest pain (anginal) and to reduce the risk of death due to heart problems in certain patients who have had a heart attack. It may also be used for other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: **WARNING:** Do not suddenly stop taking this medicine; sharp chest pain, irregular heartbeat, and sometimes heart attack may occur. The risk may be greater if you have certain types of heart disease. Your doctor should slowly lower your dose over several weeks if you need to stop taking it, even if you only take it for high blood pressure. Heart disease is common and you may not know you have it. Limit physical activity while you are lowering your dose. If new or worsened chest pain or other heart problems occur, contact your doctor right away. You may need to start taking this medicine again. Tell your doctor or dentist that you take this medicine before you receive any medical or dental care, emergency care, or surgery. Tell your doctor if you have a history of diabetes or take medicine to lower your blood sugar (eg, glyburide, insulin). This medicine may hide signs of low blood sugar such as fast heartbeat. Tell your doctor right away if you notice other signs of low blood sugar such as anxiety, chills, dizziness, drowsiness, fainting, headache, tremor, unusual sweating, vision changes, or weakness. This medicine should not usually be used by patients who have a history of certain lung or breathing problems (eg, asthma) or a certain type of adrenal gland tumor (pheochromocytoma). It may worsen these conditions. Tell your doctor if you have a history of these conditions; your doctor may need to adjust your dose or prescribe additional medicine to reduce the risk of side effects. Tell your doctor if you have a history of overactive thyroid. This medicine may hide symptoms of overactive thyroid (eg, fast heartbeat). Do not suddenly stop taking this medicine; suddenly stopping this medicine could worsen your condition. Your doctor should slowly lower your dose over several weeks if you need to stop taking it. Check with your doctor for more information. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking mifepristide. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking amiodarone, bupropion, certain HIV protease inhibitors (eg, ritonavir), certain selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), cimetidine, digoxin, difenhydramine, didoxypyrimidine, flecainide, hormonal contraceptives (eg, birth control pills), hydralazine, hydroxychloroquine, ketanserin, mefloquine, phenothiazines (eg, thioridazine), propafenone, quinzalofine (eg, alfuzosin), quinidine, terbinafine, propylthiouracil, verapamil, barbiturates (eg, phenobarbital), indomethacin, phenylpropanolamine, bupivacaine, lidocaine, or clonidine. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions including lung or breathing problems (eg, asthma, bronchitis, chronic obstructive pulmonary disease [COPD]), diabetes, low blood pressure, thyroid problems, or adrenal gland tumor (pheochromocytoma), allergies, pregnancy, or liver disease, or if you will be having surgery or receiving anesthesia. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have a very slow heartbeat (eg, bradycardia), certain types of irregular heartbeat (eg, atrioventricular [AV] block, sick sinus syndrome), moderate to severe heart failure, very low systolic blood pressure (less than 100 mm Hg), or severe blood circulation problems. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for taking this medicine provided by your doctor. TAKE THIS MEDICINE BY MOUTH WITH FOOD or immediately following a meal at the same time each day. STORE THIS MEDICINE at room temperature, between 59 and 86 degrees F (15 and 30 degrees C). Store away from heat, moisture, and light. Do not store in the bathroom. If YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

CAUTIONS: DO NOT USE THIS MEDICINE if you are allergic to any ingredient in this medicine or to another beta-blocker (eg, propranolol). IF YOU HAVE A HISTORY OF ANY SEVERE ALLERGIC REACTION, talk with your doctor. You may be at risk for an even more severe allergic reaction if you come into contact with the substance that caused your allergy. Some medicines used to treat

severe allergies may also not work as well while you are using this medicine. **LAB TESTS**, including liver and kidney function, blood pressure, and complete blood cell counts, may be performed while you use this medicine. These tests may be used to monitor your condition or check for side effects. Be sure to keep all doctor and lab appointments. **THIS MEDICINE MAY CAUSE DROWSINESS, DIZZINESS, OR LIGHTEADEDNESS.** These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. Do not drive or perform other possibly unsafe tasks until you know how you react to it. This medicine may cause dizziness, lightheadedness, or fainting; alcohol, hot weather, exercise, or fever may increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of any of these effects. **PATIENTS WHO TAKE MEDICINE FOR HIGH BLOOD PRESSURE** often feel tired or run down for a few weeks after starting treatment. Be sure to take your medicine even if you may not feel "normal." Tell your doctor if you develop any new symptoms. **DO NOT SUDDENLY STOP TAKING THIS MEDICINE** without first consulting your doctor. If your doctor decides you should no longer use this medicine, you will need to stop this medicine gradually according to your doctor's instructions. **IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK YOUR BLOOD PRESSURE** and heart rate regularly, be sure to do so. **TELL YOUR DOCTOR OR DENTIST** that you take this medicine before you receive any medical or dental care, emergency care, or surgery. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES**, either prescription or over-the-counter, check with your doctor or pharmacist. Do not take any medicines used to treat colds or congestion without first talking with your doctor or pharmacist. **FOR WOMEN:** If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. This medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby. **DIABETES PATIENTS:** this medicine may hide signs of low blood sugar, such as rapid heartbeat. Be sure to watch for other signs or low blood sugar. Low blood sugar may make you anxious, sweaty, weak, dizzy, drowsy, or faint. It may also make your vision change; give you a headache, chills, or tremors; or make you more hungry. Check blood sugar levels closely. Ask your doctor before you change the dose of your diabetes medicine.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while taking this medicine include constipation, diarrhea, dizziness, dry mouth/eyes, gas, headache, heartburn, lightheadedness, mild drowsiness, muscle aches, nausea, stomach pain, trouble sleeping, unusual tiredness or weakness, or vomiting. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience blue or unusually cold hands or feet; chest pain; fainting; hallucinations; mood or mental changes (eg, confusion, depression); pounding in the chest; severe dizziness or lightheadedness; shortness of breath; slow or irregular heartbeat; swelling of the arms, hands, and feet; vision changes; wheezing; yellowing of the skin or eyes. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include chest pain, seizures, very slow heart beat, severe dizziness, fainting, and difficult or slowed breathing.

ADDITIONAL INFORMATION: DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WICH 957918

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

NELVA BRUNSTING
13630 Pinetack, Houston, TX 770797517
(713)464-4391
RX # 1534699-03328 DATE: 04/29/11

METOPROLOL TARTRATE 50MG TABLETS
QTY: 60 1 REFILL BEFORE 04/05/12
Refill NDC: 00378-0032-10
Retail Price: \$16.66 Your Insurance Saved You: \$12.28 **\$ 4.38**

A. JAIN, MD PLAN: PADMNPD
MFG:MYLAN GROUP# CMD3896
XXX/PBP/PBP/ /PBP CLAIM REF# SLOMDCX

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024
PH: (713)722-7247 **Customer Receipt**

NELVA BRUNSTING
13630 Pinetack, Houston, TX 770797517
(713)464-4391
RX # 1534699-03328 DATE: 04/29/11

METOPROLOL TARTRATE 50MG TABLETS
QTY: 60 1 REFILL BEFORE 04/05/12
Refill NDC: 00378-0032-10
Retail Price: \$16.66 Your Insurance Saved You: \$12.28 **\$ 4.38**

A. JAIN, MD PLAN: PADMNPD
MFG:MYLAN GROUP# CMD3896
XXX/PBP/PBP/ /PBP CLAIM REF# SLOMDCX

Walgreens 12850 MEMORIAL DRIVE HOUSTON, TX 77024
PH: (713)722-7247 **Duplicate Receipt**

Pharmacy use only
METOPROLOL TARTRATE 50MG TABLETS
SUN 11:00AM 00378-0032-10
Refill CELL 138

QTY 60
20 DRAM

PINK
FRONT: m 32
XXX/PBP/PBP/ /PBP

Brnstrng004459



Cold or flu? Antibiotics aren't for you.

Antibiotics kill bacteria,
not viruses.

- Antibiotics can cure most bacterial infections, such as sore throats caused by strep and bacterial sinus infections.
- Using antibiotics for viral illness, like the common cold, will not help you feel better or prevent spreading it.

Please follow your healthcare provider's advice. And to learn more about antibiotics, visit www.cdc.gov/getsmart today.

5745



When you're sick, antibiotics aren't always the answer.

To avoid antibiotic-resistant infections and adverse drug events, avoid seeking an antibiotic prescription for colds, coughs and sniffles.

- Taking antibiotics when they are not needed causes some bacteria to become resistant to the antibiotic.
- Resistant bacteria are stronger and make future bacterial illnesses harder to treat.

To learn more, talk with your
Walgreens pharmacist today.

OPR: 2875 67 119 9963941

INFO: 0017 00378003210

5745

8
BR




PAIDMPD *1534699 0302 3 0000438 5*

NELVA BRUNSTING
13630 Pinerock
Houston, TX 770797517
(713)464-4391

- Store TT 2 of 5
- Your Insurance Saved You: \$12.28

SUN 11:00AM
\$4.38
EXPRESS PAY

04/29/11
Refill



**Personal
Prescription
Information**

LOOK INSIDE FOR IMPORTANT INFORMATION ABOUT YOUR MEDICATION.

Take advantage of this convenient service:

- **Auto Refills**
We'll automatically refill your prescription before it's due to run out. Sign up in the pharmacy.

Walgreens
The Pharmacy America Trusts • Since 1901™

Visit us online at Walgreens.com

Thank you for choosing Walgreens!

Brunsting004460